

CERTAIN SUBCLASSES OF ANALYTIC FUNCTIONS WITH VARYING ARGUMENTS

MOHAMED K. AOUF - ADELA O. MOSTAFA - EMAN A. ADWAN

In this paper, we introduce new classes $VM(\beta)$ and $VN(\beta)$ of analytic functions with varying arguments in the open unit disc $U = \{z \in \mathbb{C} : |z| < 1\}$. Some properties such as coefficient estimates, extreme points, distortion theorems for functions $f(z)$ belonging to the classes are obtained.

1. Introduction

Let A denote the class of functions of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n, \quad (1)$$

which are analytic in the open unit disc $U = \{z : |z| < 1\}$. Let $M(\beta)$ denote the subclass of A consisting of functions $f(z)$ which satisfy the inequality:

$$\Re \left\{ \frac{zf'(z)}{f(z)} \right\} < \beta, \quad (2)$$

or, equivalently,

$$\left| \frac{\frac{zf'(z)}{f(z)} - 1}{\frac{zf'(z)}{f(z)} - (2\beta - 1)} \right| < 1, \quad (3)$$

Entrato in redazione: 22 aprile 2013

AMS 2010 Subject Classification: 30C45.

Keywords: Harmonic univalent functions, Dziok- Srivastava operator, Extreme points.

for some $\beta (\beta > 1)$. Also let $N(\beta)$ denote the subclass of A consisting of functions $f(z)$ which satisfy the inequality:

$$\Re \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} < \beta, \quad (4)$$

or, equivalently,

$$\left| \frac{\frac{zf''(z)}{f'(z)}}{1 + \frac{zf''(z)}{f'(z)} - (2\beta - 1)} \right| < 1. \quad (5)$$

The classes $M(\beta)$ and $N(\beta)$ were introduced by Nishiwaki and Owa [3] (see also [1], [2], [4], [5] and [7]) and for $1 < \beta \leq \frac{4}{3}$ the classes $M(\beta)$ and $N(\beta)$ were introduced and studied by Uralegaddi et al. (see [8]). It follows from (2) and (4) we can see that (see [7])

$$f(z) \in N(\beta) \iff zf'(z) \in M(\beta).$$

Definition 1.1 ([6]). A function $f(z)$ defined by (1) is said to be in the class $V(\theta_n)$ if $f(z) \in A$ and $\arg(a_n) = \theta_n$ for all $n \geq 2$. If furthermore, there exists a real number α such that

$$\theta_n + (n-1)\alpha \equiv \pi \pmod{2\pi},$$

then $f(z)$ is said to be in the class $V(\theta_n; \alpha)$. The union of $V(\theta_n; \alpha)$ taken over all possible sequences $\{\theta_n\}$ and all possible real numbers α is denoted by V .

Denote by $VM(\beta)$ the subclass of V consisting of functions $f(z)$ in $M(\beta)$ and denote by $VN(\beta)$ the subclass of V consisting of functions $f(z)$ in $N(\beta)$.

2. Coefficient estimates

Unless otherwise mentioned, we shall assume in the reminder of this paper that $1 < \beta \leq \frac{4}{3}$.

To prove our main results we shall need the following lemmas.

Lemma 2.1 ([8]). *If $f(z) \in A$ satisfies*

$$\sum_{n=2}^{\infty} (n-\beta) |a_n| \leq \beta - 1, \quad (6)$$

then $f(z) \in M(\beta)$.

Lemma 2.2 ([8]). *If $f(z) \in A$ satisfies*

$$\sum_{n=2}^{\infty} n(n - \beta) |a_n| \leq \beta - 1, \tag{7}$$

then $f(z) \in N(\beta)$.

Theorem 2.3. *Let the function $f(z)$ be of the form (1), then $f(z)$ is in the class $VM(\beta)$ if and only if*

$$\sum_{n=2}^{\infty} (n - \beta) |a_n| \leq \beta - 1. \tag{8}$$

Proof. In view of Lemma 2.1, we need only to show the function $f(z)$ from the class $VM(\beta)$ satisfies the coefficient inequality (8). Let $f(z) \in VM(\beta)$. Then, from (1) and (3), we have

$$\left| \frac{\sum_{n=2}^{\infty} (n - 1) a_n z^{n-1}}{2(\beta - 1) + \sum_{n=2}^{\infty} (n - 2\beta + 1) a_n z^{n-1}} \right| < 1.$$

Since $f(z) \in V$, $f(z)$ lies in the class $V(\theta_n, \alpha)$ for some sequence $\{\theta_n\}$ and a real number α such that $\theta_n + (n - 1)\alpha \equiv \pi \pmod{2\pi}$ ($n \geq 2$), then setting $z = re^{i\alpha}$ in the above inequality, we get

$$\left| \frac{-\sum_{n=2}^{\infty} (n - 1) |a_n| r^{n-1}}{2(\beta - 1) - \sum_{n=2}^{\infty} (n - 2\beta + 1) |a_n| r^{n-1}} \right| < 1.$$

Since $Re\{w(z)\} < |w(z)| < 1$, we have

$$\Re \left\{ \frac{\sum_{n=2}^{\infty} (n - 1) |a_n| r^{n-1}}{2(\beta - 1) - \sum_{n=2}^{\infty} (n - 2\beta + 1) |a_n| r^{n-1}} \right\} < 1. \tag{9}$$

Hence

$$\sum_{n=2}^{\infty} (n - \beta) |a_n| r^{n-1} \leq (\beta - 1),$$

which, upon letting $r \rightarrow 1^-$, readily yields the assertion (8). This completes the proof of Theorem 2.3. □

Corollary 2.4. Let the function $f(z)$ defined by (1) be in the class $VM(\beta)$, then

$$|a_n| \leq \frac{(\beta - 1)}{(n - \beta)} \quad (n \geq 2). \quad (10)$$

The result is sharp for the function

$$f(z) = z + \frac{(\beta - 1)}{(n - \beta)} e^{i\theta_n} z^n \quad (n \geq 2). \quad (11)$$

Similarly, we can prove the following theorem for the class $VN(\beta)$.

Theorem 2.5. Let the function $f(z)$ be of the form (1), then $f(z)$ is in the class $VN(\beta)$ if and only if

$$\sum_{n=2}^{\infty} n(n - \beta) |a_n| \leq (\beta - 1). \quad (12)$$

Corollary 2.6. Let the function $f(z)$ defined by (1) be in the class $VN(\beta)$, then

$$|a_n| \leq \frac{(\beta - 1)}{n(n - \beta)} \quad (n \geq 2). \quad (13)$$

The result is sharp for the function

$$f(z) = z + \frac{(\beta - 1)}{n(n - \beta)} e^{i\theta_n} z^n \quad (n \geq 2). \quad (14)$$

3. Distortion theorem

Theorem 3.1. Let the function $f(z)$ defined by (1) be in the class $VM(\beta)$. Then

$$|z| - \frac{(\beta - 1)}{(2 - \beta)} |z|^2 \leq |f(z)| \leq |z| + \frac{(\beta - 1)}{(2 - \beta)} |z|^2, \quad (15)$$

The result is sharp.

Proof. Since

$$\Psi(n) = (n - \beta), \quad (16)$$

is an increasing function of n ($n \geq 2$), from Theorem 2.3, we have

$$(2 - \beta) \sum_{n=2}^{\infty} |a_n| \leq \sum_{n=2}^{\infty} (n - \beta) |a_n| \leq (\beta - 1),$$

that is

$$\sum_{n=2}^{\infty} |a_n| \leq \frac{(\beta - 1)}{(2 - \beta)},$$

Thus

$$\begin{aligned}
 |f(z)| &= \left| z + \sum_{n=2}^{\infty} a_n z^n \right| \leq |z| + |z|^2 \sum_{n=2}^{\infty} |a_n| \\
 &\leq |z| + \frac{(\beta - 1)}{(2 - \beta)} |z|^2.
 \end{aligned}$$

Similarly, we get

$$\begin{aligned}
 |f(z)| &\geq |z| - \sum_{n=2}^{\infty} |a_n| |z|^n \geq |z| - |z|^2 \sum_{n=2}^{\infty} |a_n| \\
 &\geq |z| - \frac{(\beta - 1)}{(2 - \beta)} |z|^2.
 \end{aligned}$$

This completes the proof of Theorem 3.1. Finally the result is sharp for the function

$$f(z) = z + \frac{(\beta - 1)}{(2 - \beta)} e^{i\theta_2} z^2 \tag{17}$$

$$z = \pm |z| e^{-i\theta_2}.$$

□

Corollary 3.2. *Under the hypotheses of Theorem 3.1, $f(z)$ is included in a disc with center at the origin and radius r_1 given by*

$$r_1 = 1 + \frac{(\beta - 1)}{(2 - \beta)}.$$

Theorem 3.3. *Let the function $f(z)$ defined by (1) belong to the class $VM(\beta)$. Then*

$$1 - \frac{2(\beta - 1)}{(2 - \beta)} |z| \leq |f'(z)| \leq 1 + \frac{2(\beta - 1)}{(2 - \beta)} |z|. \tag{18}$$

The result is sharp for the function $f(z)$ given by (17) at $z = \pm |z| e^{-i\theta_2}$.

Proof. Since $\{n\Psi(n)\}$, where $\Psi(n)$ given by (16) is increasing function of n ($n \geq 2$), then in view of Theorem 2.5, we have

$$\frac{(2 - \beta)}{2} \sum_{n=2}^{\infty} n |a_n| \leq \sum_{n=2}^{\infty} (n - \beta) |a_n| \leq (\beta - 1),$$

that is

$$\sum_{n=2}^{\infty} n |a_n| \leq \frac{2(\beta - 1)}{(2 - \beta)}.$$

Thus

$$\begin{aligned} |f'(z)| &= \left| 1 + \sum_{n=2}^{\infty} n a_n z^{n-1} \right| \leq 1 + |z| \sum_{n=2}^{\infty} n |a_n| \\ &\leq 1 + \frac{2(\beta-1)}{(2-\beta)} |z|. \end{aligned}$$

Similarly, we get

$$\begin{aligned} |f'(z)| &\geq 1 - \sum_{n=2}^{\infty} n |a_n| |z|^{n-1} \geq 1 - |z| \sum_{n=2}^{\infty} n |a_n| \\ &\geq 1 - \frac{2(\beta-1)}{(2-\beta)} |z|. \end{aligned}$$

Finally the result is sharp for the function $f(z)$ given by (17). This completes the proof of Theorem 3.3. \square

Corollary 3.4. *Let the function $f(z)$ defined by (1) be in the class $VM(\beta)$. Then $f'(z)$ is included in a disc with center at the origin and radius r_2 given by*

$$r_2 = 1 + \frac{2(\beta-1)}{(2-\beta)}.$$

Using the same technique as used in Theorems 3.1 and 3.3, we have the following theorems for functions in the class $VN(\beta)$:

Theorem 3.5. *Let the function $f(z)$ defined by (1) be in the class $VN(\beta)$. Then*

$$|z| - \frac{(\beta-1)}{2(2-\beta)} |z|^2 \leq |f(z)| \leq |z| + \frac{(\beta-1)}{2(2-\beta)} |z|^2,$$

The result is sharp for the function

$$f(z) = z + \frac{(\beta-1)}{2(2-\beta)} e^{i\theta_2} z^2 \quad (19)$$

at $z = \pm |z| e^{-i\theta_2}$.

Corollary 3.6. *Under the hypotheses of Theorem 3.5, $f(z)$ is included in a disc with center at the origin and radius r_3 given by*

$$r_3 = 1 + \frac{(\beta-1)}{2(2-\beta)}.$$

Theorem 3.7. *Let the function $f(z)$ defined by (1) belong to the class $VN(\beta)$. Then*

$$1 - \frac{(\beta - 1)}{(2 - \beta)} |z| \leq |f'(z)| \leq 1 + \frac{(\beta - 1)}{(2 - \beta)} |z|.$$

The result is sharp for the function $f(z)$ given by (19) at $z = \pm |z|e^{-i\theta_2}$.

Corollary 3.8. *Let the function $f(z)$ defined by (1) be in the class $VN(\beta)$. Then $f'(z)$ is included in a disc with center at the origin and radius r_4 given by*

$$r_4 = 1 + \frac{(\beta - 1)}{(2 - \beta)}.$$

4. Extreme points

Theorem 4.1. *Let the function $f(z)$ defined by (1) be in the class $VM(\beta)$, with $\arg(a_n) = \theta_n$ where $[\theta_n + (n - 1)\alpha] \equiv \pi \pmod{2\pi}$. Define $f_1(z) = z$ and*

$$f_n(z) = z + \frac{(\beta - 1)}{(n - \beta)} e^{i\theta_n} z^n. \tag{20}$$

Then $f(z)$ is in the class $VM(\beta)$ if and only if it can be expressed in the form

$$f(z) = \sum_{n=1}^{\infty} \mu_n f_n(z), \tag{21}$$

where $\mu_n \geq 0$ ($n \geq 1$) and $\sum_{n=1}^{\infty} \mu_n = 1$.

Proof. If $f(z) = \sum_{n=1}^{\infty} \mu_n f_n(z)$ with $\sum_{n=1}^{\infty} \mu_n = 1$ and $\mu_n \geq 0$, then

$$\begin{aligned} & \sum_{n=2}^{\infty} (n - \beta) \frac{(\beta - 1)}{(n - \beta)} \mu_n \\ &= \sum_{n=2}^{\infty} (\beta - 1) \mu_n = (\beta - 1)(1 - \mu_1) \leq (\beta - 1). \end{aligned}$$

Hence $f(z) \in VM(\beta)$.

Conversely, let the function $f(z)$ defined by (1) belongs to the class $VM(\beta)$, define

$$\mu_n = \frac{(n - \beta)}{(\beta - 1)} |a_n|, \tag{22}$$

and

$$\mu_1 = 1 - \sum_{n=2}^{\infty} \mu_n.$$

From Theorem 2.3, $\sum_{n=1}^{\infty} \mu_n \leq 1$ and so $\mu_n \geq 0$. Since $\mu_n f_n(z) = \mu_n z + a_n z^n$, then

$$\sum_{n=1}^{\infty} \mu_n f_n(z) = z + \sum_{n=2}^{\infty} a_n z^n = f(z).$$

This completes the proof of Theorem 4.1. □

Similarly, we can prove the following theorem for the class $VN(\beta)$.

Theorem 4.2. *Let the function $f(z)$ defined by (1) be in the class $VN(\beta)$, with $\arg(a_n) = \theta_n$ where $[\theta_n + (n-1)\alpha] \equiv \pi \pmod{2\pi}$. Define $f_1(z) = z$ and*

$$f_n(z) = z + \frac{(\beta-1)}{n(n-\beta)} e^{i\theta_n} z^n. \quad (23)$$

Then $f(z)$ is in the class $VN(\beta)$ if and only if it can be expressed in the form

$$f(z) = \sum_{n=1}^{\infty} \mu_n f_n(z), \quad (24)$$

where $\mu_n \geq 0$ ($n \geq 1$) and $\sum_{n=1}^{\infty} \mu_n = 1$.

REFERENCES

- [1] M. K. Aouf - A. Shamandy - A. O. Mostafa - E. A. Adwan, *Subordination results for certain class of analytic functions defined by convolution*, Rend. del Circolo Mat. di Palermo 60 (2011), 255–262.
- [2] M. K. Aouf - A. Shamandy - A. O. Mostafa - E. A. Adwan, *Subordination theorem of analytic functions defined by convolution*, Complex Anal. and Oper. Theory (To appear).
- [3] J. Nishiwaki - S. Owa, *Coefficient inequalities for certain analytic functions*, Internat. J. Math. Math. Sci. 29 (5) (2002), 285–290.
- [4] S. Owa - J. Nishiwaki, *Coefficient estimates for certain classes of analytic functions*, J. Inequal. Pure Appl. Math. 3 (5) (2002), Art. 72, 1–12.
- [5] S. Owa - H. M. Srivastava, *Some generalized convolution properties associated with certain subclasses of analytic functions*, J. Inequal. Pure Appl. Math. 3 (3) (2002), Art.42, 1–27.
- [6] H. Silverman, *Univalent functions with varying arguments*, Houston J. Math. 7 (1981), 283–287.

- [7] H. M. Srivastava - A. A. Attiya, *Some subordination results associated with certain subclasses of analytic functions*, J. Inequal. Pure Appl. Math. 5 (4) (2004), Art.82, 1–14.
- [8] B. A. Uralegaddi - M. D. Ganigi - S. M. Sarangi, *Univalent functions with positive coefficients*, Tamkang J. Math. 25 (3) (1994), 225–230.

MOHAMED K. AOUF

*Department of Mathematics, Faculty of Science,
Mansoura University, Mansoura 35516, Egypt
e-mail: mkaouf127@yahoo.com*

ADELA O. MOSTAFA

*Department of Mathematics, Faculty of Science,
Mansoura University, Mansoura 35516, Egypt
e-mail: adelaeg254@yahoo.com*

EMAN A. ADWAN

*Department of Mathematics, Faculty of Science,
Mansoura University, Mansoura 35516, Egypt
e-mail: eman.a2009@yahoo.com*