

ON STABILITY FOR NONLINEAR IMPLICIT FRACTIONAL DIFFERENTIAL EQUATIONS

MOUFFAK BENCHOHRA - JAMAL E. LAZREG

The purpose of this paper is to establish some types of Ulam stability: Ulam-Hyers stability, generalized Ulam-Hyers stability, Ulam-Hyers-Rassias stability and generalized Ulam-Hyers-Rassias stability for a class of implicit fractional-order differential equation.

1. Introduction

Fractional calculus is a generalization of ordinary differentiation and integration to arbitrary order (non-integer). See, for example, the books such as [2–4, 7, 27, 31, 37] and the articles [9, 10], and references therein.

In recent years, fractional differential equations arise naturally in various fields such as rheology, fractals, chaotic dynamics, modeling and control theory, signal processing, bioengineering and biomedical applications, etc; Fractional derivatives provide an excellent instrument for the description of memory and hereditary properties of various materials and processes. We refer the reader, for example, to the books such as [8, 23, 25, 39], and references therein.

“Under what conditions does there exist an additive mapping near an approximately additive mapping?” This is the stability problem of functional equation (of group homomorphisms) which was raised by Ulam in 1940 in a talk

Entrato in redazione: 3 giugno 2014

AMS 2010 Subject Classification: 26A33, 34A08.

Keywords: Initial value problem, Caputo’s fractional derivative, implicit fractional differential equations, fractional integral, existence, Gronwall’s lemma, fixed point, Ulam-Hyers stability, Ulam-Hyers-Rassias stability.

given at Wisconsin University [40, 41]. In 1941, Hyers [17] gave the first answer to the question of Ulam (for the additive mapping) in the case Banach spaces. Between 1982 and 1998 Rassias established the Hyers-Ulam stability of linear and nonlinear mappings. Jung [19, 20] investigated in 1998, the Hyers-Ulam stability of more general mappings on restricted domains. Obloza [26], in 1997, is the first author who has investigated the Hyers-Ulam stability of linear differential equations. After, many articles and books on this subject have been published in order to generalize the results of Hyers in many directions. For more detailed definitions of the Hyers-Ulam stability and the generalized Hyers-Ulam stability, we refer the reader to the papers [1, 5, 6, 16, 18, 21, 22, 24, 32, 36, 43–45] and the books [12, 33, 34].

Integer order implicit differential equations of arbitrary orders have been considered extensively in the literature; see for instance [11, 13–15, 28–30, 35, 38, 46].

The purpose of this paper, is to establish four types of Ulam stability : Ulam-Hyers stability, generalized Ulam-Hyers stability, Ulam-Hyers-Rassias stability and generalized Ulam-Hyers-Rassias stability for the equation, for the following initial value problem for implicit fractional-order differential equation

$${}^c D^\alpha y(t) = f(t, y(t), {}^c D^\alpha y(t)), \quad \forall t \in J, \quad 0 < \alpha \leq 1, \quad (1)$$

$$y(0) = y_0, \quad (2)$$

where ${}^c D^\alpha$ is the Caputo fractional derivative, $f : J \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is a given function space, $y_0 \in \mathbb{R}$ and $J = [0, T]$, $T > 0$.

This paper initiates the study of Ulam stability for such class of problems.

2. Preliminaries

Definition 2.1 ([31]). The fractional (arbitrary) order integral of the function $h \in L^1([0, T], \mathbb{R}_+)$ of order $\alpha \in \mathbb{R}_+$ is defined by

$$I^\alpha h(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} h(s) ds,$$

where Γ is the Euler gamma function defined by $\Gamma(\alpha) = \int_0^\infty t^{\alpha-1} e^{-t} dt$, $\alpha > 0$.

Definition 2.2 ([23]). For a function h given on the interval $[0, T]$, the Caputo fractional-order derivative of order α of h , is defined by

$$({}^c D^\alpha h)(t) = \frac{1}{\Gamma(n-\alpha)} \int_0^t (t-s)^{n-\alpha-1} h^{(n)}(s) ds,$$

where $n = [\alpha] + 1$ and $[\alpha]$ denotes the integer part of the real number α .

Lemma 2.3 ([23]). *Let $\alpha \geq 0$ and $n = [\alpha] + 1$. Then*

$$I^\alpha({}^c D^\alpha f(t)) = f(t) - \sum_{k=0}^{n-1} \frac{f^{(k)}(0)}{k!} t^k.$$

We state the following generalization of Gronwall's lemma for singular kernels.

Lemma 2.4 ([42]). *Let $v : [0, T] \rightarrow [0, +\infty)$ be a real function and $w(\cdot)$ is a nonnegative, locally integrable function on $[0, T]$ and there are constants $a > 0$ and $0 < \alpha < 1$ such that*

$$v(t) \leq w(t) + a \int_0^t (t-s)^{-\alpha} v(s) ds,$$

Then, there exists a constant $K = K(\alpha)$ such that

$$v(t) \leq w(t) + Ka \int_0^t (t-s)^{-\alpha} w(s) ds, \text{ for every } t \in [0, T].$$

We adopt the definitions in Rus [36]: Ulam-Hyers stability, generalized Ulam-Hyers stability, Ulam-Hyers-Rassias stability and generalized Ulam-Hyers-Rassias stability for the equation, for the implicit fractional-order differential equation (1).

Definition 2.5. The equation (1) is *Ulam-Hyers stable* if there exists a real number $c_f > 0$ such that for each $\varepsilon > 0$ and for each solution $z \in C^1(J, \mathbb{R})$ of the inequality

$$|{}^c D^\alpha z(t) - f(t, z(t), {}^c D^\alpha z(t))| \leq \varepsilon, \quad t \in J, \quad (3)$$

there exists a solution $y \in C^1(J, \mathbb{R})$ of equation (1) with

$$|z(t) - y(t)| \leq c_f \varepsilon, \quad t \in J.$$

Definition 2.6. The equation (1) is *generalised Ulam-Hyers stable* if there exists $\psi_f \in C(\mathbb{R}_+, \mathbb{R}_+)$, $\psi_f(0) = 0$, such that for each solution $z \in C^1(J, \mathbb{R})$ of the inequality (3) there exists a solution $y \in C^1(J, \mathbb{R})$ of the equation (1) with

$$|z(t) - y(t)| \leq \psi_f(\varepsilon), \quad t \in J.$$

Definition 2.7. The equation (1) is *Ulam-Hyers-Rassias stable* with respect to $\varphi \in C(J, \mathbb{R}_+)$ if there exists a real number $c_f > 0$ such that for each $\varepsilon > 0$ and for each solution $z \in C^1(J, \mathbb{R})$ of the inequality

$$|{}^c D^\alpha z(t) - f(t, z(t), {}^c D^\alpha z(t))| \leq \varepsilon \varphi(t), \quad t \in J, \quad (4)$$

there exists a solution $y \in C^1(J, \mathbb{R})$ of equation (1) with

$$|z(t) - y(t)| \leq c_f \varepsilon \varphi(t), \quad t \in J.$$

Definition 2.8. The equation (1) is *generalised Ulam-Hyers-Rassias stable* with respect to $\varphi \in C(J, \mathbb{R}_+)$ if there exists a real number $c_{f,\varphi} > 0$ such that for each solution $z \in C^1(J, \mathbb{R})$ of the inequality

$$|{}^c D^\alpha z(t) - f(t, z(t), {}^c D^\alpha z(t))| \leq \varphi(t), \quad t \in J, \quad (5)$$

there exists a solution $y \in C^1(J, \mathbb{R})$ of equation (1) with

$$|z(t) - y(t)| \leq c_{f,\varphi} \varphi(t), \quad t \in J.$$

Remark 2.9. A function $z \in C^1(J, \mathbb{R})$ is a solution of of the inequality (3) if and only if there exists a function $g \in C(J, \mathbb{R})$ (which depend on y) such that

$$(i) \quad |g(t)| \leq \varepsilon, \quad \forall t \in J.$$

$$(ii) \quad {}^c D^\alpha z(t) = f(t, z(t), {}^c D^\alpha z(t)) + g(t), \quad t \in J.$$

Remark 2.10. Clearly,

$$(i) \quad \text{Definition 2.5} \implies \text{Definition 2.6.}$$

$$(ii) \quad \text{Definition 2.7} \implies \text{Definition 2.8.}$$

Remark 2.11. A solution of the implicit differential inequation (3) with fractional order is called an fractional ε -solution of the implicit fractional differential equation (1).

So, the Ulam stabilities of the implicit differential equations with fractional order are some special types of data dependence of the solutions of fractional implicit differential equations.

3. Existence and Ulam-Hyers Stability

Definition 3.1. A function $u \in C^1(J)$ is said to be a solution of the problem (1) – (2) is u satisfied equation (1) and condition (2) on J .

Lemma 3.2. Let a function $f(t, u, v) : J \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ be continuous. Then the problem (1) – (2) is equivalent to the problem

$$y(t) = y_0 + I^\alpha g(t), \quad (6)$$

where $g \in C(J, \mathbb{R})$ satisfies the functional equation

$$g(t) = f(t, y_0 + I^\alpha g(t), g(t)).$$

Proof. If ${}^c D^\alpha y(t) = g(t)$ then $I^\alpha {}^c D^\alpha y(t) = I^\alpha g(t)$. So we obtain $y(t) = y_0 + I^\alpha g(t)$. \square

Lemma 3.3 ([9]). *Assume*

(H1) *The function $f : J \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous.*

(H2) *There exist constants $K > 0$ and $0 < L < 1$ such that*

$$|f(t, u, v) - f(t, \bar{u}, \bar{v})| \leq K|u - \bar{u}| + L|v - \bar{v}| \text{ for any } u, v, \bar{u}, \bar{v} \in \mathbb{R} \text{ and } t \in J.$$

If

$$\frac{KT^\alpha}{(1-L)\Gamma(\alpha+1)} < 1, \quad (7)$$

then there exists a unique solution for the IVP (1) – (2) on J .

Theorem 3.4. *Assume that the assumptions (H1), (H2) and (7) hold. Then the equation (1) is Ulam-Hyers stable.*

Proof. Let $z \in C(J, \mathbb{R})$ be a solution of the inequation (3), i.e.

$$|{}^c D^\alpha z(t) - f(t, z(t), {}^c D^\alpha z(t))| \leq \varepsilon, \quad t \in J. \quad (8)$$

Let us denote by $y \in C(J, \mathbb{R})$ the unique solution of the Cauchy problem

$$\begin{aligned} {}^c D^\alpha y(t) &= f(t, y(t), {}^c D^\alpha y(t)), \quad \forall t \in J, \quad 0 < \alpha \leq 1, \\ y(0) &= z(0). \end{aligned}$$

By using Lemma 3.2, we have

$$y(t) = z(0) + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} g_y(s) ds,$$

where $g_y \in C(J, \mathbb{R})$ satisfies the functional equation

$$g_y(t) = f(t, y(0) + I^\alpha g_y(t), g_y(t)).$$

But, by integration of the formula (8) we obtain

$$\begin{aligned} \left| z(t) - z(0) - \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} g_z(s) ds \right| &\leq \frac{\varepsilon t^\alpha}{\Gamma(\alpha+1)} \\ &\leq \frac{\varepsilon T^\alpha}{\Gamma(\alpha+1)}, \end{aligned} \quad (9)$$

where $g_z \in C(J, \mathbb{R})$ satisfies the functional equation

$$g_z(t) = f(t, z(0) + I^\alpha g_z(t), g_z(t)).$$

On the other hand, we have, for each $t \in J$

$$\begin{aligned} |z(t) - y(t)| &= \left| z(t) - z(0) - \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} g_y(s) ds \right| \\ &= \left| z(t) - z(0) - \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} g_z(s) ds \right. \\ &\quad \left. + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} (g_z(s) - g_y(s)) ds \right| \\ &\leq \left| z(t) - z(0) - \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} g_z(s) ds \right| \\ &\quad + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} |g_z(s) - g_y(s)| ds, \end{aligned} \quad (10)$$

where

$$g_y(t) = f(t, y(t), g_y(t)),$$

and

$$g_z(t) = f(t, z(t), g_z(t)).$$

By (H2), we have, for each $t \in J$

$$\begin{aligned} |g_z(t) - g_y(t)| &= |f(t, z(t), g_z(t)) - f(t, y(t), g_y(t))| \\ &\leq K|z(t) - y(t)| + L|g_z(t) - g_y(t)|. \end{aligned}$$

Then

$$|g_z(t) - g_y(t)| \leq \frac{K}{1-L} |z(t) - y(t)|. \quad (11)$$

Thus, by (9), (10), and (11) we get

$$|z(t) - y(t)| \leq \frac{\varepsilon T^\alpha}{\Gamma(\alpha+1)} + \frac{K}{(1-L)\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} |z(s) - y(s)| ds.$$

Then Lemma 2.4 implies that for each $t \in J$

$$|z(t) - y(t)| \leq \frac{\varepsilon T^\alpha}{\Gamma(\alpha+1)} \left[1 + \frac{\gamma K T^\alpha}{(1-L)\Gamma(\alpha+1)} \right] := c\varepsilon, \quad (12)$$

where $\gamma = \gamma(\alpha)$ is a constant. So, the equation (1) is Ulam-Hyers stable. This completes the proof. \square

By putting $\psi(\varepsilon) = c\varepsilon$, $\psi(0) = 0$ yields that the equation (1) is generalized Ulam-Hyers stable.

4. Ulam-Hyers-Rassias Stability.

Theorem 4.1. Assume (H1), (H2) and

(H3) The function $\varphi \in C(J, \mathbb{R}_+)$ is increasing and there exists $\lambda_\varphi > 0$ such that, for each $t \in J$, we have

$$I^\alpha \varphi(t) \leq \lambda_\varphi \varphi(t).$$

Then the equation (1) is Ulam-Hyers-Rassias stable with respect to φ .

Proof. Let $z \in C(J, \mathbb{R})$ be a solution of the inequation (4), i.e.

$$|{}^c D^\alpha z(t) - f(t, z(t), {}^c D^\alpha z(t))| \leq \varepsilon \varphi(t), \quad t \in J, \quad \varepsilon > 0. \quad (13)$$

Let us denote by $y \in C(J, \mathbb{R})$ the unique solution of the Cauchy problem

$$\begin{aligned} {}^c D^\alpha y(t) &= f(t, y(t), {}^c D^\alpha y(t)), \quad \forall t \in J, \quad 0 < \alpha \leq 1, \\ y(0) &= z(0). \end{aligned}$$

By using Lemma 3.2, we have

$$y(t) = z(0) + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} g_y(s) ds,$$

where $g_y \in C(J, \mathbb{R})$ satisfies the functional equation

$$g_y(t) = f(t, y(0) + I^\alpha g_y(t), g_y(t)).$$

But, by integration of the formula (13) and by (H3), we obtain

$$\begin{aligned} \left| z(t) - z(0) - \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} g_z(s) ds \right| &\leq \frac{\varepsilon}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} \varphi(s) ds \\ &\leq \varepsilon \lambda_\varphi \varphi(t), \end{aligned} \quad (14)$$

where $g_z \in C(J, \mathbb{R})$ satisfies the functional equation

$$g_z(t) = f(t, z(0) + I^\alpha g_z(t), g_z(t)).$$

On the other hand, we have, for each $t \in J$

$$\begin{aligned} |z(t) - y(t)| &= \left| z(t) - z(0) - \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} g_y(s) ds \right| \\ &= \left| z(t) - z(0) - \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} g_z(s) ds \right. \\ &\quad \left. + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} (g_z(s) - g_y(s)) ds \right| \\ &\leq \left| z(t) - z(0) - \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} g_z(s) ds \right| \\ &\quad + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} |g_z(s) - g_y(s)| ds, \end{aligned} \quad (15)$$

where

$$g_y(t) = f(t, y(t), g_y(t)),$$

and

$$g_z(t) = f(t, z(t), g_z(t)).$$

By (H2), we have

$$\begin{aligned} |g_z(t) - g_y(t)| &= |f(t, z(t), g_z(t)) - f(t, y(t), g_y(t))| \\ &\leq K|z(t) - y(t)| + L|g_z(t) - g_y(t)|. \end{aligned}$$

Then

$$|g_z(t) - g_y(t)| \leq \frac{K}{1-L}|z(t) - y(t)|. \quad (16)$$

Thus, by (14), (15), and (16)

$$|z(t) - y(t)| \leq \varepsilon \lambda_\varphi \varphi(t) + \frac{K}{(1-L)\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} |z(s) - y(s)| ds.$$

Then Lemma 2.4 implies that, for each $t \in J$

$$|z(t) - y(t)| \leq \varepsilon \lambda_\varphi \varphi(t) + \frac{\gamma_1 \varepsilon \lambda_\varphi K}{(1-L)\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} \varphi(s) ds, \quad (17)$$

where constant $\gamma_1 = \gamma_1(\alpha)$ is a constant.

Thus, by (H3) and (17), we obtain

$$\begin{aligned} |z(t) - y(t)| &\leq \varepsilon \lambda_\varphi \varphi(t) + \frac{\gamma_1 K \varepsilon \lambda_\varphi^2 \varphi(t)}{(1-L)} \\ &= \left(1 + \frac{\gamma_1 K \lambda_\varphi}{1-L}\right) \lambda_\varphi \varepsilon \varphi(t). \end{aligned}$$

Then, for each $t \in J$

$$|z(t) - y(t)| \leq \left[\left(1 + \frac{\gamma_1 K \lambda_\varphi}{1-L}\right) \lambda_\varphi \right] \varepsilon \varphi(t) := c \varepsilon \varphi(t). \quad (18)$$

So, the equation (1) is Ulam-Hyers-Rassias stable. This completes the proof. \square

5. Examples

Example 5.1. Consider the following Cauchy problem

$${}^c D^{\frac{1}{2}} y(t) = \frac{1}{100} (t \cos y(t) - y(t) \sin(t)) + \frac{|{}^c D^{\frac{1}{2}} y(t)|}{50 + |{}^c D^{\frac{1}{2}} y(t)|}, \quad \forall t \in [0, 1], \quad (19)$$

$$y(0) = 1. \quad (20)$$

Set

$$f(t, u, v) = \frac{1}{100} (t \cos u - u \sin(t)) + \frac{v}{50 + v}, \quad t \in [0, 1], \quad u, v \in \mathbb{R}.$$

Clearly, the function f is jointly continuous.

For any $u, v, \bar{u}, \bar{v} \in \mathbb{R}$ and $t \in [0, 1]$:

$$\begin{aligned} |f(t, u, v) - f(t, \bar{u}, \bar{v})| &\leq \frac{1}{100} |t| |\cos u - \cos \bar{u}| \\ &\quad + \frac{1}{100} |\sin t| |u - \bar{u}| + \frac{50|v - \bar{v}|}{(50 + v)(50 + \bar{v})} \\ &\leq \frac{1}{100} |u - \bar{u}| + \frac{1}{100} |u - \bar{u}| + \frac{1}{50} |v - \bar{v}| \\ &\leq \frac{1}{50} |u - \bar{u}| + \frac{1}{50} |v - \bar{v}|. \end{aligned}$$

Hence condition (H2) is satisfied with $K = L = \frac{1}{50}$.

Thus condition

$$\frac{KT^\alpha}{(1-L)\Gamma(\alpha+1)} = \frac{\frac{1}{50}}{(1-\frac{1}{50})\Gamma(\frac{3}{2})} = \frac{2}{49\sqrt{\pi}} < 1,$$

is satisfied. It follows from Theorem 3.3 that the problem (19)-(20) has a unique solution on J , and from Theorem 3.4, equation (19) is Ulam-Hyers stable.

Example 5.2. Consider the following Cauchy problem

$${}^c D^{\frac{1}{2}} y(t) = \frac{2 + |y(t)| + |{}^c D^{\frac{1}{2}} y(t)|}{120e^{t+10}(1 + |y(t)| + |{}^c D^{\frac{1}{2}} y(t)|)}, \quad \forall t \in [0, 1], \quad (21)$$

$$y(0) = 1. \quad (22)$$

Set

$$f(t, u, v) = \frac{2 + |u| + |v|}{120e^{t+10}(1 + |u| + |v|)}, \quad t \in [0, 1], \quad u, v \in \mathbb{R}.$$

Clearly, the function f is jointly continuous.

For any $u, v, \bar{u}, \bar{v} \in \mathbb{R}$ and $t \in [0, 1]$

$$|f(t, u, v) - f(t, \bar{u}, \bar{v})| \leq \frac{1}{120e^{10}}(|u - \bar{u}| + |v - \bar{v}|).$$

Hence condition (H2) is satisfied with $K = L = \frac{1}{120e^{10}}$.

Let $\varphi(t) = t^2$. We have

$$I^\alpha \varphi(t) \leq \frac{2}{\Gamma(\frac{7}{2})} t^2 := \lambda_\varphi \varphi(t).$$

Thus condition (H3) is satisfied with $\varphi(t) = t^2$ and $\lambda_\varphi = \frac{2}{\Gamma(\frac{7}{2})} = \frac{16}{15\sqrt{\pi}}$. It follows from Theorem 3.3 that the problem (21)-(22) as a unique solution on J , and from Theorem 4.1 equation (21) is Ulam-Hyers-Rassias stable.

REFERENCES

- [1] S. Abbas - M. Benchohra, *On the generalized Ulam-Hyers-Rassias stability for Darboux problem for partial fractional implicit differential equations*, Appl. Math. E-Notes 14 (2014), 20–28.
- [2] S. Abbas - M. Benchohra - G. M. N'Guérékata, *Topics in Fractional Differential Equations*, Springer-Verlag, New York, 2012.
- [3] S. Abbas - M. Benchohra - G. M. N'Guérékata, *Advanced Fractional Differential and Integral Equations*, Nova Science Publishers, New York, 2014.
- [4] G. A. Anastassiou, *Advances on Fractional Inequalities*, Springer, New York Dordrecht Heidelberg London, 2011.
- [5] C. Alsina - R. Ger, *On some inequalities and stability results related to the exponential function*, J. Inequal. Appl. 2 (1998), 373–380.
- [6] T. Aoki, *On the stability of the linear transformation in Banach spaces*, J. Math. Soc. Japan 2 (1950), 64–66.
- [7] D. Baleanu - K. Diethelm - E. Scalas - J. J. Trujillo, *Fractional Calculus Models and Numerical Methods*, World Scientific Publishing, New York, 2012.
- [8] D. Baleanu - Z. B. Güvenç - J. A. T. Machado, *New Trends in Nanotechnology and Fractional Calculus Applications*, Springer, New York, 2010.
- [9] M. Benchohra - J. E. Lazreg, *Nonlinear fractional implicit differential equations*, Commun. Appl. Anal. 17 (2013), 471–482.
- [10] M. Benchohra - J. E. Lazreg, *Existence and uniqueness results for nonlinear implicit fractional differential equations with boundary conditions*, Rom. J. Math. Comput. Sc. 4 (1) (2014), 60–72.

- [11] R. Bielawski - L. G \check{z} orniewicz: *A fixed point index approach to some differential equations*, in Proc. Conf. Topological Fixed Point Theory and Appl. (Boju Jiang, Ed.), Lect. Not. Math. 1411, pp. 9–14, Springer-Verlag, New York, 1989.
- [12] Y. J. Cho - Th. M. Rassias - R. Saadati, *Stability of Functional Equations in Random Normed Spaces*, Science-Business Media, Springer 52, 2013.
- [13] L. H. Erbe - W. Krawcewicz - T. Kaczynski, *Solvability of two-point boundary value problems for systems of nonlinear differential equations of the form $y'' = g(t, y, y', y'')$* , Rocky Mountain J. Math. 20 (1990), 899–907.
- [14] M. Feckan, *Existence results for implicit differential equations*, Math. Slovaca 48 (1998), 35–42.
- [15] M. Frigon - T. Kaczynski, *Boundary value problems for systems of implicit differential equations*, J. Math. Anal. Appl. 179 (1993), 317–326.
- [16] P. Gavruta, *A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings*, J. Math. Anal. Appl. 184 (1994), 431–436.
- [17] D. H. Hyers, *On the stability of the linear functional equation*, Natl. Acad.Sci. U.S.A. 27 (1941), 222–224.
- [18] R. W. Ibrahim, *Stability for univalent solutions of complex fractional differential equations*, Proc. Pakistan Acad. Sci. 49 (3) (2012), 227–232.
- [19] S. M. Jung, *On the Hyers-Ulam stability of the functional equations that have the quadratic property*, J. Math. Anal. Appl. 222 (1998), 126–137.
- [20] S. M. Jung, *Hyers-Ulam stability of linear differential equations of first order II*, Appl. Math. Lett. 19 (2006), 854–858.
- [21] K. W. Jun - H. M. Kim, *On the stability of an n-dimensional quadratic and additive functional equation*, Math. Inequal. Appl. 19 (9) (2006), 854–858.
- [22] S. M. Jung - K. S. Lee, *Hyers-Ulam stability of first order linear partial differential equations with constant coefficients*, Math. Inequal. Appl. 10 (2007), 261–266.
- [23] A. A. Kilbas - Hari M. Srivastava - Juan J. Trujillo, *Theory and Applications of Fractional Differential Equations*, North-Holland Mathematics Studies, 204, Elsevier Science B.V., Amsterdam, 2006.
- [24] G. H. Kim, *On the stability of functional equations with square-symmetric operation*, Math. Inequal. Appl. 17 (4) (2001), 257–266.
- [25] V. Lakshmikantham - S. Leela - J. Vasundhara, *Theory of Fractional Dynamic Systems*, Cambridge Academic Publishers, Cambridge, 2009.
- [26] M. Obloza, *Hyers stability of the linear differential equation*, Rocznik Nauk-Dydakt. Prace Mat. 13 (1993), 259–270.
- [27] M. D. Ortigueira, *Fractional Calculus for Scientists and Engineers*, Lecture Notes in Electrical Engineering, 84, Springer, Dordrecht, 2011.
- [28] W. V. Petryshyn, *Solvability of various boundary value problems for the equation $x'' = f(t, x, x', x'') - y$* , Pacific J. Math. 122 (1986), 169–195.

- [29] W. V. Petryshyn - Z. S. Yu, *On the solvability of an equation describing the periodic motions of a satellite in its elliptic orbit*, Nonlinear Anal., Th. Meth. Appl. 9 (1985), 969–975.
- [30] W. V. Petryshyn - Z. S. Yu, *Solvability of Neumann boundary problems for nonlinear second-order odes which need not be solvable for the highest-order derivative*, J. Math. Anal. Appl. 91 (1983), 244–253.
- [31] I. Podlubny, *Fractional Differential Equations*, Academic Press, San Diego, 1999.
- [32] Th. M. Rassias, *On the stability of the linear mapping in Banach spaces*, Proc. Amer. Math. Soc. 72 (1978), 297–300.
- [33] J. M. Rassias, *Functional Equations, Difference Inequalities and Ulam Stability Notions (F.U.N.)*, Nova Science Publishers, Inc. New York, 2010.
- [34] Th. M. Rassias - J. Brzdek, *Functional Equations in Mathematical Analysis*, Springer 86, New York Dordrecht Heidelberg London 2012.
- [35] B. Ricceri, *On the Cauchy problem for the differential equation $f(t, x, x', \dots, x^{(k)}) = 0$* , Glasgow Math. J. 33 (1991), 343–348.
- [36] I. A. Rus, *Ulam stabilities of ordinary differential equations in a Banach space*, Carpathian J. Math. 26 (2010), 103–107.
- [37] S. G. Samko - A. A. Kilbas - O. I. Marichev, *Fractional Integrals and Derivatives. Theory and Applications*, Gordon and Breach, Yverdon, 1993.
- [38] K. R. Schneider, *Existence and approximation results to the Cauchy problem for a class of differential-algebraic equations*, Zeit. Anal. Anwen. 10 (1991), 375–384.
- [39] V. E. Tarasov, *Fractional Dynamics: Application of Fractional Calculus to Dynamics of particles, Fields and Media*, Springer, Heidelberg; Higher Education Press, Beijing, 2010.
- [40] S. M. Ulam, *Problems in Modern Mathematics*, Chapter 6, John Wiley and Sons, New York, USA, 1940.
- [41] S. M. Ulam, *A Collection of Mathematical Problems*, Interscience, New York, 1960.
- [42] H. Ye - J. Gao - Y. Ding, *A generalized Gronwall inequality and its application to a fractional differential equation*, J. Math. Anal. Appl. 328 (2007), 1075–1081.
- [43] J. Wang - M. Fečkan - Y. Zhou, *Ulam's type stability of impulsive ordinary differential equations*, J. Math. Anal. Appl. 395 (2012), 258–264.
- [44] J. Wang - L. Lv - Y. Zhou, *Ulam stability and data dependence for fractional differential equations with Caputo derivative*, Electron. J. Qual. Theory Differ. Equat. 63 (2011), 1–10.
- [45] J. Wang - Y. Zhang, *Existence and stability of solutions to nonlinear impulsive differential equations in β -normed spaces*, Electron. J. Differential Equations 83 (2014), 1–10.
- [46] J. R. L. Webb - S. C. Welsh, *Existence and uniqueness of initial value problems for a class of second-order differential equations*, J. Differential Equations 82 (1989), 314–321.

MOUFFAK BENCHOHRA
Laboratory of Mathematics
University of Sidi Bel-Abbès
P.O. Box 89, Sidi Bel Abbès 22000, Algeria
and
Department of Mathematics
King Abdulaziz University
Faculty of Science
P.O. Box 80203, Jeddah 21589, Saudi Arabia
e-mail: benchohra@univ-sba.dz

JAMAL E. LAZREG
Laboratory of Mathematics
University of Sidi Bel Abbès
P.O. Box 89, Sidi Bel-Abbès 22000, Algeria
e-mail: lazregjamal@yahoo.fr