

CALDERÓN'S REPRODUCING FORMULAS FOR THE SPHERICAL MEAN L^2 -MULTIPLIER OPERATORS

KHALED HLEILI

First we study the spherical mean L^2 -multiplier operators on $[0, +\infty[\times \mathbb{R}^n$.
Next, we give for these operators Calderón's reproducing formulas and
best approximation formulas.

1. Introduction

In the Euclidean case the multiplier operator T_m associated with a bounded function m on \mathbb{R}^n is defined by $\widehat{T_m f} = m \widehat{f}$, where \widehat{f} denotes the classical Fourier transform. Many authors [5, 9, 24] have been interested to extend the L^p Fourier-multipliers on several hypergroups and to show similarly its L^p -boundedness. Recently, these operators are studied in [25] where the author established some applications (Calderón's reproducing formulas, best approximation formulas and extremal functions...).

The spherical mean operator \mathcal{R} is defined, for a function f on $\mathbb{R} \times \mathbb{R}^n$, even with respect to the first variable [15], by

$$\mathcal{R}(f)(r, x) = \int_{S^n} f(r\eta, x + r\xi) d\sigma_n(\eta, \xi), \quad (r, x) \in \mathbb{R} \times \mathbb{R}^n,$$

Entrato in redazione: 30 novembre 2016

AMS 2010 Subject Classification: 43A32, 42B10

Keywords: Spherical mean operator, L^2 -multiplier operators, Calderón's reproducing formulas, best approximation formulas.

where S^n is the unit sphere of $\mathbb{R} \times \mathbb{R}^n$ and $d\sigma_n$ is the surface measure on S^n normalized to have total measure one.

The dual of the spherical mean operator ${}^t\mathcal{R}$ is defined by

$${}^t\mathcal{R}(g)(r, x) = \frac{1}{(2\pi)^{\frac{n}{2}}} \int_{\mathbb{R}^n} g(\sqrt{r^2 + |x - y|^2}, y) dy.$$

The spherical mean operator \mathcal{R} and its dual have many important physical applications, namely in image processing of so-called synthetic aperture radar (SAR) data [6, 7, 23, 28], or in the linearized inverse scattering problem in acoustics [4].

The Fourier transform \mathcal{F} associated with the spherical mean operator is defined for every integrable function f on $[0, +\infty[\times \mathbb{R}^n$ with respect to the measure $d\nu_{n+1}$, by

$$\forall (s, y) \in \Upsilon, \mathcal{F}(f)(s, y) = \int_0^{+\infty} \int_{\mathbb{R}^n} f(r, x) \mathcal{R}(\cos(s \cdot) e^{-i(y|\cdot|)})(r, x) d\nu_{n+1}(r, x),$$

where $d\nu_{n+1}$ is the measure defined on $[0, +\infty[\times \mathbb{R}^n$ by

$$d\nu_{n+1}(r, x) = \frac{r^n}{2^{\frac{n-1}{2}} \Gamma(\frac{n+1}{2})} dr \otimes \frac{dx}{(2\pi)^{\frac{n}{2}}},$$

$\|\cdot\|_{p, \nu_{n+1}}$ its norm, and Υ is the set given by

$$\Upsilon = \mathbb{R} \times \mathbb{R}^n \cup \{(ir, x), (r, x) \in \mathbb{R} \times \mathbb{R}^n, |r| \leq |x|\}. \quad (1.1)$$

Many harmonic analysis results related to the Fourier transform \mathcal{F} have already been proved by Dziri, Jlassi, Nessibi, Rachdi and Trimche [3, 8, 15, 18] or also by Peng and Zhao [17, 30]. Recently, Baccar, Omri and Rachdi [2] studied the generalized Fock spaces associated with the spherical mean operator \mathcal{R} , and Msehli, Rachdi and Omri [13, 14, 16] established several uncertainty principles for the Fourier transform \mathcal{F} .

Let m be a function in the Lebesgue space $L^2(d\nu_{n+1})$. We define the spherical mean L^2 -multiplier operators on $[0, +\infty[\times \mathbb{R}^n$, for regular functions

$$T_{m, \varepsilon} f = \mathcal{F}^{-1}((m_\varepsilon \circ \theta) \mathcal{F}(f)), \quad \varepsilon > 0,$$

where m_ε is the function given by

$$m_\varepsilon(r, x) = m(\varepsilon r, \varepsilon x), \quad (1.2)$$

and θ is the bijective function, defined on the set

$$\Upsilon_+ = [0, +\infty[\times \mathbb{R}^n \cup \{(is, y) ; (s, y) \in [0, +\infty[\times \mathbb{R}^n ; s \leq |y|\}$$

by,

$$\theta(s, y) = (\sqrt{s^2 + |y|^2}, y). \tag{1.3}$$

Our purpose in this work is to study the multiplier $T_{m,\varepsilon}$, for which we shall prove an analogue of the Calderón's reproducing formulas by using the theory of the Fourier transform \mathcal{F} and the convolution product $*$.

Next, we use the theory of reproducing kernels to give best approximation of these operators and a Calderón's reproducing formula of the associated extremal function. This paper is organized as follows, in the second section we recall some harmonic analysis results related to the spherical mean operator \mathcal{R} and its associated Fourier transform \mathcal{F} .

In the third section we study the spherical mean L^2 -multiplier operators $T_{m,\varepsilon}$, and for these operators we establish Calderón's reproducing formulas.

The last section of this paper is devoted to giving best approximation for every function $m \in L^\infty(d\nu_{n+1})$ of the operators $T_{m,\varepsilon}$.

2. The spherical mean operator

In [15], Nessibi, Rachdi and Trimèche showed that for every $(\mu, \lambda) \in \mathbb{C} \times \mathbb{C}^n$, the function $\varphi_{(\mu,\lambda)}$ defined on $\mathbb{R} \times \mathbb{R}^n$ by

$$\varphi_{(\mu,\lambda)}(r, x) = \mathcal{R} \left(\cos(\mu \cdot) e^{-i\langle \lambda, \cdot \rangle} \right) (r, x), \tag{2.1}$$

is the unique infinitely differentiable function on $\mathbb{R} \times \mathbb{R}^n$, even with respect to the first variable, satisfying the following system

$$\begin{cases} \frac{\partial u}{\partial x_j}(r, x_1, \dots, x_n) = -i\lambda_j u(r, x_1, \dots, x_n), & 1 \leq j \leq n, \\ \ell_{\frac{n-1}{2}} u(r, x_1, \dots, x_n) - \Delta u(r, x_1, \dots, x_n) = -\mu^2 u(r, x_1, \dots, x_n), \\ u(0, \dots, 0) = 1, \\ \frac{\partial u}{\partial r}(0, x_1, \dots, x_n) = 0, & (x_1, \dots, x_n) \in \mathbb{R}^n, \end{cases}$$

where $\ell_{\frac{n-1}{2}}$ is the Bessel operator, defined by $\ell_{\frac{n-1}{2}} = \frac{\partial^2}{\partial r^2} + \frac{n}{r} \frac{\partial}{\partial r}$, and Δ denotes the usual Laplacian operator defined by $\Delta = \sum_{j=1}^n \frac{\partial^2}{\partial x_j^2}$. The authors proved also

that the eigenfunction $\varphi_{(\mu,\lambda)}$ defined by relation (2.1), is explicitly given by

$$\forall (r,x) \in \mathbb{R} \times \mathbb{R}^n, \quad \varphi_{(\mu,\lambda)}(r,x) = j_{\frac{n-1}{2}}(r\sqrt{\mu^2 + |\lambda|^2})e^{-i\langle \lambda, x \rangle}, \quad (2.2)$$

where $j_{\frac{n-1}{2}}$ is the modified Bessel function defined by

$$j_{\frac{n-1}{2}}(z) = 2^{\frac{n-1}{2}} \Gamma\left(\frac{n+1}{2}\right) \frac{J_{\frac{n-1}{2}}(z)}{z^{\frac{n-1}{2}}} = \Gamma\left(\frac{n+1}{2}\right) \sum_{k=0}^{+\infty} \frac{(-1)^k}{k! \Gamma\left(\frac{n+1}{2} + k\right)} \left(\frac{z}{2}\right)^{2k}, \quad z \in \mathbb{C},$$

and $J_{\frac{n-1}{2}}$ is the Bessel function of the first kind and index $\frac{n-1}{2}$ (see [1, 11] and [29]).

The modified Bessel function $j_{\frac{n-1}{2}}$ has the following integral representation

$$\forall z \in \mathbb{C}, \quad j_{\frac{n-1}{2}}(z) = \frac{2\Gamma\left(\frac{n+1}{2}\right)}{\sqrt{\pi}\Gamma\left(\frac{n}{2}\right)} \int_0^1 (1-t^2)^{\frac{n}{2}-1} \cos(zt) dt. \quad (2.3)$$

Relation (2.3) shows in particular that, for every $z \in \mathbb{C}$ and for every $k \in \mathbb{N}$, we have

$$\left| j_{\frac{n-1}{2}}^{(k)}(z) \right| \leq e^{|\operatorname{Im}(z)|}.$$

From the properties of the modified Bessel function $j_{\frac{n-1}{2}}$, we deduce that the eigenfunction $\varphi_{(\mu,\lambda)}$ is bounded on $\mathbb{R} \times \mathbb{R}^n$ if, and only if, (μ, λ) belongs to the set Υ given by relation (1.1), and in this case

$$\sup_{(r,x) \in \mathbb{R} \times \mathbb{R}^n} \left| \varphi_{(\mu,\lambda)}(r,x) \right| = 1. \quad (2.4)$$

In the following we shall define the translation operators, the convolution product and the Fourier transform \mathcal{F} associated with the operator \mathcal{B} . For this we denote by

- \mathcal{B}_{Υ_+} the σ -algebra defined on Υ_+ by,

$$\mathcal{B}_{\Upsilon_+} = \{ \theta^{-1}(B), B \in \mathcal{B}_{\text{Bor}}([0, +\infty[\times \mathbb{R}^n) \},$$

where θ is the function, given by relation (1.3).

- γ_{n+1} the measure defined on \mathcal{B}_{Υ_+} by, $\gamma_{n+1}(B) = \nu_{n+1}(\theta(B))$.
- $L^p(d\gamma_{n+1})$, $p \in [1, +\infty]$ the Lebesgue space of measurable functions f on Υ_+ , such that $\|f\|_{p, \gamma_{n+1}} < +\infty$.

We have the following properties (see [15] and [26])

i) For every nonnegative measurable function g on Υ_+ , we have

$$\int \int_{\Upsilon_+} g(\mu, \lambda) d\gamma_{n+1}(\mu, \lambda) = \frac{1}{2^{\frac{n-1}{2}} \Gamma(\frac{n+1}{2}) (2\pi)^{\frac{n}{2}}} \left(\int_0^{+\infty} \int_{\mathbb{R}^n} g(\mu, \lambda) (\mu^2 + |\lambda|^2)^{\frac{n-1}{2}} \mu d\mu d\lambda + \int_{\mathbb{R}^n} \int_0^{|\lambda|} g(i\mu, \lambda) (|\lambda|^2 - \mu^2)^{\frac{n-1}{2}} \mu d\mu d\lambda \right).$$

ii) For every nonnegative measurable function f on $[0, +\infty[\times \mathbb{R}^n$ (respectively integrable on $[0, +\infty[\times \mathbb{R}^n$ with respect to the measure $d\nu_{n+1}$), $f \circ \theta$ is a measurable nonnegative function on Υ_+ , (respectively integrable on Υ_+ with respect to the measure $d\gamma_{n+1}$) and we have

$$\int \int_{\Upsilon_+} (f \circ \theta)(\mu, \lambda) d\gamma_{n+1}(\mu, \lambda) = \int_0^{+\infty} \int_{\mathbb{R}^n} f(r, x) d\nu_{n+1}(r, x).$$

Moreover, the function f belongs to $L^p(d\nu_{n+1})$, $p \in [1, +\infty]$ if and only if $f \circ \theta$ belongs to $L^p(d\gamma_{n+1})$ and we have

$$\|f\|_{p, \nu_{n+1}} = \|f \circ \theta\|_{p, \gamma_{n+1}}. \tag{2.5}$$

According to Rachdi, Nessibi and Trimèche (see [15, 26] and [27]), we have the following definition and properties for the translation operator associated with the spherical mean operator

Definition 2.1. i) For every $(r, x) \in [0, +\infty[\times \mathbb{R}^n$, the translation operator $\mathcal{T}_{(r,x)}$ associated with the spherical mean operator is defined on $L^p(d\nu_{n+1})$, $p \in [1, +\infty]$, by

$$\mathcal{T}_{(r,x)}(f)(s, y) = \frac{\Gamma(\frac{n+1}{2})}{\sqrt{\pi} \Gamma(\frac{n}{2})} \int_0^\pi f(\sqrt{r^2 + s^2 + 2rs \cos \theta}, x + y) \sin^{n-1}(\theta) d\theta.$$

ii) The convolution product of measurable functions f and g on $[0, +\infty[\times \mathbb{R}^n$, is defined by

$$\forall (r, x) \in [0, +\infty[\times \mathbb{R}^n; f * g(r, x) = \int_0^{+\infty} \int_{\mathbb{R}^n} \mathcal{T}_{(r,x)}(f)(s, -y) g(s, y) d\nu_{n+1}(s, y),$$

whenever the integral of the right-hand side is defined.

For every $(r, x) \in]0, +\infty[\times \mathbb{R}^n$, and by a standard change of variables, we have

$$\forall (s, y) \in]0, +\infty[\times \mathbb{R}^n, \mathcal{T}_{(r,x)}(f)(s, y) = \frac{1}{2^{\frac{n-1}{2}} \Gamma(\frac{n+1}{2})} \int_0^{+\infty} f(t, x + y) \mathcal{W}_n(r, s, t) t^n dt,$$

where the kernel \mathcal{W}_n , is given by

$$\mathcal{W}_n(r, s, t) = \frac{\Gamma(\frac{n+1}{2})^2}{2^{\frac{n-3}{2}}\Gamma(\frac{n}{2})\sqrt{\pi}} \frac{((r+s)^2 - t^2)^{\frac{n}{2}-1} (t^2 - (r-s)^2)^{\frac{n}{2}-1}}{(rst)^{n-1}} \mathbf{1}_{|r-s|, r+s}(t).$$

Also, the coming properties are satisfied

• For every $f \in L^p(d\nu_{n+1})$, $p \in [1, +\infty]$, and $(r, x) \in [0, +\infty[\times \mathbb{R}^n$, the function $\mathcal{T}_{(r,x)}(f)$ belongs to $L^p(d\nu_{n+1})$ and we have

$$\|\mathcal{T}_{(r,x)}(f)\|_{p, \nu_{n+1}} \leq \|f\|_{p, \nu_{n+1}}. \quad (2.6)$$

• Let $p, q, r \in [1, +\infty]$ such that $\frac{1}{p} + \frac{1}{q} = 1 + \frac{1}{r}$. Then for every $f \in L^p(d\nu_{n+1})$ and $g \in L^q(d\nu_{n+1})$, the function $f * g$ belongs to the space $L^r(d\nu_{n+1})$, and we have the following Young's inequality

$$\|f * g\|_{r, \nu_{n+1}} \leq \|f\|_{p, \nu_{n+1}} \|g\|_{q, \nu_{n+1}}.$$

In the following, we shall define the Fourier transform \mathcal{F} connected with the spherical mean operator, and we recall some of its properties that we need in the next sections.

Definition 2.2. The Fourier transform \mathcal{F} associated with the spherical mean operator is defined on $L^1(d\nu_{n+1})$ by [15]

$$\forall (\mu, \lambda) \in \Upsilon; \mathcal{F}(f)(\mu, \lambda) = \int_0^{+\infty} \int_{\mathbb{R}^n} f(r, x) \varphi_{(\mu, \lambda)}(r, x) d\nu_{n+1}(r, x),$$

where $\varphi_{(\mu, \lambda)}$ is the eigenfunction given by relation (2.2), and Υ is the set defined by relation (1.1).

Then, according to [15], we have
For every $f, g \in L^1(d\nu_{n+1})$,

$$\mathcal{F}(f * g) = \mathcal{F}(f)\mathcal{F}(g),$$

and $(\mu, \lambda) \in \Upsilon$

$$\mathcal{F}(\mathcal{T}_{(r,-x)}(f))(\mu, \lambda) = \varphi_{(\mu, \lambda)}(r, x) \mathcal{F}(f)(\mu, \lambda). \quad (2.7)$$

Moreover, relation (2.4) implies that the Fourier transform \mathcal{F} is a bounded linear operator from $L^1(d\nu_{n+1})$ into $L^\infty(d\gamma_{n+1})$, and that for every $f \in L^1(d\nu_{n+1})$, we have

$$\|\mathcal{F}(f)\|_{\infty, \gamma_{n+1}} \leq \|f\|_{1, \nu_{n+1}}. \quad (2.8)$$

For every positive real number ε and for every $m \in L^p(d\nu_{n+1})$, $p \in [1, +\infty[$, the function m_ε defined by relation (1.2), belongs to $L^p(d\nu_{n+1})$ and we have

$$\|m_\varepsilon\|_{p, \nu_{n+1}} = \frac{1}{\varepsilon^{\frac{2n+1}{p}}} \|m\|_{p, \nu_{n+1}}. \quad (2.9)$$

In [15], Rachdi, Nessibi and Trimèche, established the following inversion formula and Plancherel theorem for the Fourier transform \mathcal{F} .

Theorem 2.3 (Inversion formula). *Let $f \in L^1(d\nu_{n+1})$ such that $\mathcal{F}(f) \in L^1(d\gamma_{n+1})$, then for almost every $(r, x) \in \mathbb{R} \times \mathbb{R}^n$*

$$f(r, x) = \int \int_{\Gamma_+} \mathcal{F}(f)(\mu, \lambda) \overline{\varphi_{(\mu, \lambda)}(r, x)} d\gamma_{n+1}(\mu, \lambda).$$

Theorem 2.4 (Plancherel theorem). *The Fourier transform \mathcal{F} can be extended to an isometric isomorphism from $L^2(d\nu_{n+1})$ onto $L^2(d\gamma_{n+1})$. In particular, for every $f \in L^2(d\nu_{n+1})$*

$$\|\mathcal{F}(f)\|_{2, \gamma_{n+1}} = \|f\|_{2, \nu_{n+1}}.$$

Corollary 2.5. *For all functions f and g in $L^2(d\nu_{n+1})$, we have*

$$\int_0^{+\infty} \int_{\mathbb{R}^n} f(r, x) \overline{g(r, x)} d\nu_{n+1}(r, x) = \int \int_{\Gamma_+} \mathcal{F}(f)(\mu, \lambda) \overline{\mathcal{F}(g)(\mu, \lambda)} d\gamma_{n+1}(\mu, \lambda).$$

Remark 2.6. (i) For every $f, g \in L^2(d\nu_{n+1})$; the function $f * g$ belongs to the space $C_{e,0}(\mathbb{R} \times \mathbb{R}^n)$ consisting of continuous functions h on $\mathbb{R} \times \mathbb{R}^n$, even with respect to the first variable and such that $\lim_{r^2+|x|^2 \rightarrow +\infty} h(r, x) = 0$.

Moreover,

$$f * g = \mathcal{F}^{-1}(\mathcal{F}(f)\mathcal{F}(g)), \quad (2.10)$$

where \mathcal{F}^{-1} is the mapping defined on $L^1(d\gamma_{n+1})$ by

$$\mathcal{F}^{-1}(g)(r, x) = \int \int_{\Gamma_+} g(\mu, \lambda) \overline{\varphi_{(\mu, \lambda)}(r, x)} d\gamma_{n+1}(\mu, \lambda).$$

(ii) Let $f, g \in L^2(d\nu_{n+1})$, the function $f * g$ belongs to $L^2(d\nu_{n+1})$ if and only if $\mathcal{F}(f)\mathcal{F}(g)$ belongs to $L^2(d\gamma_{n+1})$, and we have

$$\mathcal{F}(f * g) = \mathcal{F}(f)\mathcal{F}(g).$$

(iii) Let $f, g \in L^2(d\nu_{n+1})$, then

$$\|\mathcal{F}(f)\mathcal{F}(g)\|_{2, \gamma_{n+1}} = \|f * g\|_{2, \nu_{n+1}}. \quad (2.11)$$

(iv) For every $g \in L^1(d\gamma_{n+1})$, $\mathcal{F}^{-1}(g)$ belongs to $L^\infty(d\nu_{n+1})$, and we have

$$\|\mathcal{F}^{-1}(g)\|_{\infty, \nu_{n+1}} \leq \|g\|_{1, \gamma_{n+1}}.$$

3. The Spherical mean L^2 -multiplier operators

In this section we study the spherical mean L^2 -multiplier operators on $[0, +\infty[\times \mathbb{R}^n$ and for these operators we establish Calderón's reproducing formulas.

Definition 3.1. Let m be a function in $L^2(d\nu_{n+1})$ and let ε be a positive real number. The spherical mean L^2 -multiplier operators is defined for regular functions f on $[0, +\infty[\times \mathbb{R}^n$, by

$$\forall (r, x) \in [0, +\infty[\times \mathbb{R}^n, \quad T_{m,\varepsilon}f(r, x) = \mathcal{F}^{-1}((m_\varepsilon \circ \theta)\mathcal{F}(f))(r, x), \quad (3.1)$$

where m_ε is the function given by relation (1.2) and θ is the function defined by (1.3).

Proposition 3.2. (i) For every $m \in L^2(d\nu_{n+1})$, and $f \in L^1(d\nu_{n+1})$, the function $T_{m,\varepsilon}f$ belongs to $L^2(d\nu_{n+1})$, and we have

$$\|T_{m,\varepsilon}f\|_{2,\nu_{n+1}} \leq \frac{1}{\varepsilon^{\frac{2n+1}{2}}} \|m\|_{2,\nu_{n+1}} \|f\|_{1,\nu_{n+1}}.$$

(ii) For every $m \in L^2(d\nu_{n+1})$, and $f \in L^2(d\nu_{n+1})$, then $T_{m,\varepsilon}f \in L^\infty(d\nu_{n+1})$, and we have

$$T_{m,\varepsilon}f(r, x) = \int \int_{\Upsilon_+} (m_\varepsilon \circ \theta)(\mu, \lambda) \mathcal{F}(f)(\mu, \lambda) \overline{\varphi_{(\mu,\lambda)}(r, x)} d\gamma_{n+1}(\mu, \lambda),$$

and

$$\|T_{m,\varepsilon}f\|_{\infty,\nu_{n+1}} \leq \frac{1}{\varepsilon^{\frac{2n+1}{2}}} \|m\|_{2,\nu_{n+1}} \|f\|_{2,\nu_{n+1}}.$$

(iii) For every $m \in L^\infty(d\nu_{n+1})$, and $f \in L^2(d\nu_{n+1})$, the function $T_{m,\varepsilon}f$ belongs to $L^2(d\nu_{n+1})$, and we have

$$\|T_{m,\varepsilon}f\|_{2,\nu_{n+1}} \leq \|m\|_{\infty,\nu_{n+1}} \|f\|_{2,\nu_{n+1}}.$$

Proof. (i) From relations (2.5), (2.8), (3.1), and Theorem 2.4, the function $T_{m,\varepsilon}$ belongs to $L^2(d\nu_{n+1})$, and we have

$$\begin{aligned} \|\mathcal{F}(T_{m,\varepsilon}f)\|_{2,\gamma_{n+1}} &= \|(m_\varepsilon \circ \theta)\mathcal{F}(f)\|_{2,\gamma_{n+1}} \\ &\leq \|(m_\varepsilon \circ \theta)\|_{2,\gamma_{n+1}} \|\mathcal{F}(f)\|_{\infty,\gamma_{n+1}} \\ &\leq \|m_\varepsilon\|_{2,\nu_{n+1}} \|f\|_{1,\nu_{n+1}}. \end{aligned}$$

Then, the result follows from (2.9), and Theorem 2.4.

(ii) Using (2.5), (3.1), and Remark 2.6 (iv), for every $m \in L^2(d\nu_{n+1})$, and $f \in L^2(d\nu_{n+1})$, the function $T_{m,\varepsilon}f \in L^\infty(d\nu_{n+1})$, and we have

$$\|T_{m,\varepsilon}f\|_{\infty,\nu_{n+1}} \leq \|(m_\varepsilon \circ \theta)\mathcal{F}(f)\|_{1,\gamma_{n+1}}.$$

From Hölder's inequality, relation (2.9), and Theorem 2.4, we obtain

$$\begin{aligned} \|T_{m,\varepsilon}f\|_{\infty, \nu_{n+1}} &\leq \| (m_\varepsilon \circ \theta) \|_{2, \gamma_{n+1}} \| \mathcal{F}(f) \|_{2, \gamma_{n+1}} \\ &= \| m_\varepsilon \|_{2, \nu_{n+1}} \| f \|_{2, \nu_{n+1}}, \\ &= \frac{1}{\varepsilon^{\frac{2n+1}{2}}} \| m \|_{2, \nu_{n+1}} \| f \|_{2, \nu_{n+1}}. \end{aligned}$$

Part (iii) follows from (2.5), (3.1), and Theorem 2.4. \square

Remark 3.3. According to relation (2.10), for every $m \in L^2(d\nu_{n+1})$ and $f \in L^2(d\nu_{n+1})$, we can write the spherical mean L^2 - multiplier as

$$\forall (r, x) \in [0, +\infty[\times \mathbb{R}^n, \quad T_{m,\varepsilon}f(r, x) = \mathcal{F}^{-1}(m_\varepsilon \circ \theta) * f(r, x). \quad (3.2)$$

Theorem 3.4. Let m be a function in $L^2(d\nu_{n+1})$, satisfying the admissibility condition

$$\int_0^{+\infty} |m_\varepsilon \circ \theta(\mu, \lambda)|^2 \frac{d\varepsilon}{\varepsilon} = 1, \quad (\mu, \lambda) \in \Upsilon. \quad (3.3)$$

(i) **Plancherel formula:** For every $f \in L^2(d\nu_{n+1})$, we have

$$\|f\|_{2, \nu_{n+1}}^2 = \int_0^{+\infty} \|T_{m,\varepsilon}f\|_{2, \nu_{n+1}}^2 \frac{d\varepsilon}{\varepsilon}.$$

(ii) **First Calderón's formula:** Let f be a function in $L^1(d\nu_{n+1})$, such that $\mathcal{F}(f)$ in $L^1(d\gamma_{n+1})$, we have

$$f(r, x) = \int_0^{+\infty} (T_{m,\varepsilon}f * \mathcal{F}^{-1}(\overline{m_\varepsilon \circ \theta}))(r, x) \frac{d\varepsilon}{\varepsilon}, \quad \text{a.e. } (r, x) \in [0, +\infty[\times \mathbb{R}^n.$$

Proof. (i) From relations (2.11) and (3.2), we have

$$\begin{aligned} \int_0^{+\infty} \|T_{m,\varepsilon}f\|_{2, \nu_{n+1}}^2 \frac{d\varepsilon}{\varepsilon} &= \int_0^{+\infty} \int_0^{+\infty} \int_{\mathbb{R}^n} |(\mathcal{F}^{-1}(m_\varepsilon \circ \theta) * f)(r, x)|^2 d\nu_{n+1}(r, x) \frac{d\varepsilon}{\varepsilon} \\ &= \int_0^{+\infty} \int \int_{\Upsilon_+} |m_\varepsilon \circ \theta(r, x) \mathcal{F}(f)(r, x)|^2 d\gamma_{n+1}(r, x) \frac{d\varepsilon}{\varepsilon} \\ &= \int \int_{\Upsilon_+} |\mathcal{F}(f)(r, x)|^2 \left(\int_0^{+\infty} |m_\varepsilon \circ \theta(r, x)|^2 \frac{d\varepsilon}{\varepsilon} \right) d\gamma_{n+1}(r, x). \end{aligned}$$

The result follows from Theorem 2.4, and (3.3).

(ii) Let f in $L^1(d\nu_{n+1})$. According to Proposition 3.2 (i), relation (2.6), and Corollary 2.5, we have

$$\begin{aligned} &\int_0^{+\infty} (T_{m,\varepsilon}f * \mathcal{F}^{-1}(\overline{m_\varepsilon \circ \theta}))(r, x) \frac{d\varepsilon}{\varepsilon} \\ &= \int_0^{+\infty} \left[\int_0^{+\infty} \int_{\mathbb{R}^n} T_{m,\varepsilon}f(s, y) \overline{\mathcal{T}_{(r,-x)}(\mathcal{F}^{-1}(m_\varepsilon \circ \theta))(s, y)} d\nu_{n+1}(s, y) \right] \frac{d\varepsilon}{\varepsilon} \\ &= \int_0^{+\infty} \left[\int \int_{\Upsilon_+} \mathcal{F}(T_{m,\varepsilon}f)(s, y) \overline{\mathcal{F}(\mathcal{T}_{(r,-x)}(\mathcal{F}^{-1}(m_\varepsilon \circ \theta)))(s, y)} d\gamma_{n+1}(s, y) \right] \frac{d\varepsilon}{\varepsilon}. \end{aligned}$$

Using (2.7), we obtain

$$\begin{aligned} & \int_0^{+\infty} (T_{m,\varepsilon} f * \mathcal{F}^{-1}(\overline{m_\varepsilon \circ \theta})) (r, x) \frac{d\varepsilon}{\varepsilon} \\ &= \int_0^{+\infty} \left[\iint_{\Upsilon_+} \mathcal{F}(f)(s, y) \overline{\varphi_{(s,y)}(r, x)} |m_\varepsilon \circ \theta(s, y)|^2 d\gamma_{n+1}(s, y) \right] \frac{d\varepsilon}{\varepsilon}. \end{aligned}$$

Since,

$$\begin{aligned} & \int_0^{+\infty} \left[\iint_{\Upsilon_+} \left| \mathcal{F}(f)(s, y) \overline{\varphi_{(s,y)}(r, x)} \right| |m_\varepsilon \circ \theta(s, y)|^2 d\gamma_{n+1}(s, y) \right] \frac{d\varepsilon}{\varepsilon} \\ & \leq \int \int_{\Upsilon_+} |\mathcal{F}(f)(s, y)| d\gamma_{n+1}(s, y). \end{aligned}$$

Then, the result follows from Fubini's theorem, relation (3.3), and Theorem 2.3. \square

Lemma 3.5. Let $m \in L^2(d\nu_{n+1}) \cap L^\infty(d\nu_{n+1})$, satisfy the admissibility condition (3.3). For every $0 < \xi < \delta < \infty$, the function

$$\mathcal{K}_{\xi, \delta}(\mu, \lambda) = \int_\xi^\delta |m_\varepsilon \circ \theta(\mu, \lambda)|^2 \frac{d\varepsilon}{\varepsilon},$$

belongs to $L^2(d\gamma_{n+1})$, and we have

$$\|\mathcal{K}_{\xi, \delta}\|_{2, \gamma_{n+1}}^2 \leq \ln\left(\frac{\delta}{\xi}\right) \frac{\xi^{-(2n+1)} - \delta^{-(2n+1)}}{2n+1} \|m\|_{2, \nu_{n+1}}^2 \|m\|_{\infty, \nu_{n+1}}^2.$$

Proof. Using Hölder's inequality for the measure $\frac{d\varepsilon}{\varepsilon}$, we get for every $(\mu, \lambda) \in \Upsilon$

$$|\mathcal{K}_{\xi, \delta}(\mu, \lambda)|^2 \leq \ln\left(\frac{\delta}{\xi}\right) \int_\xi^\delta |m_\varepsilon \circ \theta(\mu, \lambda)|^4 \frac{d\varepsilon}{\varepsilon}.$$

From (2.5), and (2.9), we obtain

$$\begin{aligned} \|\mathcal{K}_{\xi, \delta}\|_{2, \gamma_{n+1}}^2 & \leq \ln\left(\frac{\delta}{\xi}\right) \int_\xi^\delta \left[\iint_{\Upsilon_+} |m_\varepsilon \circ \theta(\mu, \lambda)|^4 d\gamma_{n+1}(\mu, \lambda) \right] \frac{d\varepsilon}{\varepsilon} \\ & \leq \ln\left(\frac{\delta}{\xi}\right) \frac{\xi^{-(2n+1)} - \delta^{-(2n+1)}}{2n+1} \|m\|_{2, \nu_{n+1}}^2 \|m\|_{\infty, \nu_{n+1}}^2 < \infty. \end{aligned}$$

\square

Theorem 3.6. Second Calderón's formula. *Let $m \in L^2(dv_{n+1}) \cap L^\infty(dv_{n+1})$, satisfy the admissibility condition (3.3). Then for every $f \in L^2(dv_{n+1})$ and $0 < \xi < \delta < \infty$, the function*

$$f^{\xi, \delta}(r, x) = \int_{\xi}^{\delta} (T_{m, \varepsilon} f * \mathcal{F}^{-1}(\overline{m_{\varepsilon} \circ \theta})) (r, x) \frac{d\varepsilon}{\varepsilon},$$

belongs to $L^2(dv_{n+1})$ and satisfies

$$\lim_{(\xi, \delta) \rightarrow (0^+, +\infty)} \|f^{\xi, \delta} - f\|_{2, v_{n+1}} = 0. \quad (3.4)$$

Proof. From Proposition 3.2 (iii), (2.6), (2.7) and Corollary 2.5, we have

$$\begin{aligned} f^{\xi, \delta}(r, x) &= \int_{\xi}^{\delta} \left[\int_0^{+\infty} \int_{\mathbb{R}^n} T_{m, \varepsilon} f(s, y) \overline{\mathcal{T}_{(r, -x)}(\mathcal{F}^{-1}(m_{\varepsilon} \circ \theta))(s, y)} dv_{n+1}(s, y) \right] \frac{d\varepsilon}{\varepsilon} \\ &= \int_{\xi}^{\delta} \left[\iint_{\Upsilon_+} \mathcal{F}(f)(s, y) \overline{\varphi_{(s, y)}(r, x)} |m_{\varepsilon} \circ \theta(s, y)|^2 d\gamma_{n+1}(s, y) \right] \frac{d\varepsilon}{\varepsilon}. \end{aligned}$$

By Fubini-Tonnelli's theorem, Hölder's inequality, relation (2.4) and Lemma 3.5, we get

$$\begin{aligned} &\int_{\xi}^{\delta} \left[\iint_{\Upsilon_+} \left| \mathcal{F}(f)(s, y) \overline{\varphi_{(s, y)}(r, x)} \right| |m_{\varepsilon} \circ \theta(s, y)|^2 d\gamma_{n+1}(s, y) \right] \frac{d\varepsilon}{\varepsilon} \\ &\leq \int \int_{\Upsilon_+} |\mathcal{F}(f)(s, y)| \mathcal{K}_{\xi, \delta}(s, y) d\gamma_{n+1}(s, y) \\ &\leq \sqrt{\ln\left(\frac{\delta}{\xi}\right) \frac{\xi^{-(2n+1)} - \delta^{-(2n+1)}}{2n+1}} \|f\|_{2, v_{n+1}} \|m\|_{2, v_{n+1}} \|m\|_{\infty, v_{n+1}} < \infty. \end{aligned}$$

Then, from Fubini's theorem and Theorem 2.3, we obtain

$$\begin{aligned} f^{\xi, \delta}(r, x) &= \int \int_{\Upsilon_+} \mathcal{F}(f)(s, y) \overline{\varphi_{(s, y)}(r, x)} \mathcal{K}_{\xi, \delta}(s, y) d\gamma_{n+1}(s, y) \\ &= \mathcal{F}^{-1}(\mathcal{F}(f) \mathcal{K}_{\xi, \delta})(r, x). \end{aligned}$$

On the other hand, from relation (3.3), the function $\mathcal{K}_{\xi, \delta}$ belongs to $L^\infty(d\gamma_{n+1})$, from this fact and Theorem 2.4, the function $f^{\xi, \delta} \in L^2(dv_{n+1})$, and we have

$$\mathcal{F}(f^{\xi, \delta}) = \mathcal{F}(f) \mathcal{K}_{\xi, \delta}.$$

Using the previous result and Theorem 2.4, we get

$$\|f^{\xi, \delta} - f\|_{2, v_{n+1}}^2 = \int \int_{\Upsilon_+} |\mathcal{F}(f)(\mu, \lambda)|^2 (\mathcal{K}_{\xi, \delta}(\mu, \lambda) - 1)^2 d\gamma_{n+1}(\mu, \lambda).$$

The relation (3.4) follows from $\lim_{(\xi, \delta) \rightarrow (0^+, +\infty)} \mathcal{K}_{\xi, \delta}(\mu, \lambda) = 1$, and the dominated convergence theorem. \square

4. The extremal function related to spherical mean L^2 -multiplier operators

In this section, by using the theory of extremal function and reproducing Kernel of Hilbert space [19–22], we study the extremal function associated to the spherical mean L^2 -multiplier operators. The main result of this section can be stated as follows.

Definition 4.1. Let σ be a positive function on Υ satisfying :

$$\sigma(\mu, \lambda) \geq 1, \quad (\mu, \lambda) \in \Upsilon, \quad (4.1)$$

and

$$\frac{1}{\sigma} \in L^1(d\gamma_{n+1}). \quad (4.2)$$

We define the space $\Omega_\sigma([0, +\infty[\times \mathbb{R}^n)$, by

$$\Omega_\sigma([0, +\infty[\times \mathbb{R}^n) = \{f \in L^2(d\nu_{n+1}), \sqrt{\sigma} \mathcal{F}(f) \in L^2(d\gamma_{n+1})\}.$$

The space $\Omega_\sigma([0, +\infty[\times \mathbb{R}^n)$ provided with inner product

$$\langle f, g \rangle_\sigma = \int \int_{\Upsilon_+} \sigma(\mu, \lambda) \mathcal{F}(f)(\mu, \lambda) \overline{\mathcal{F}(g)(\mu, \lambda)} d\gamma_{n+1}(\mu, \lambda),$$

and the norm $\|f\|_\sigma = \sqrt{\langle f, f \rangle_\sigma}$ is a Hilbert space.

Proposition 4.2. Let $m \in L^\infty(d\nu_{n+1})$. For every $f \in \Omega_\sigma([0, +\infty[\times \mathbb{R}^n)$, the operators $T_{m,\varepsilon}$ are bounded linear operators from $\Omega_\sigma([0, +\infty[\times \mathbb{R}^n)$ into $L^2(d\nu_{n+1})$, and we have

$$\|T_{m,\varepsilon} f\|_{2,\nu_{n+1}} \leq \|m\|_{\infty,\nu_{n+1}} \|f\|_\sigma.$$

Proof. Let $f \in \Omega_\sigma([0, +\infty[\times \mathbb{R}^n)$. According to Proposition 3.2(iii), the operator $T_{m,\varepsilon} f$ belongs to $L^2(d\nu_{n+1})$, and

$$\|T_{m,\varepsilon} f\|_{2,\nu_{n+1}} \leq \|m\|_{\infty,\nu_{n+1}} \|f\|_{2,\nu_{n+1}}.$$

By relation (4.1), we have $\|f\|_\sigma^2 \geq \int \int_{\Upsilon_+} |\mathcal{F}(f)(\mu, \lambda)|^2 d\gamma_{n+1}(\mu, \lambda)$, which gives the result. \square

Definition 4.3. Let $\rho > 0$, and let $m \in L^\infty(d\nu_{n+1})$, we denote by $\langle \cdot, \cdot \rangle_{\sigma,\rho}$ the inner product defined on the space $\Omega_\sigma([0, +\infty[\times \mathbb{R}^n)$ by

$$\langle f, g \rangle_{\sigma,\rho} = \int \int_{\Upsilon_+} (\rho \sigma(\mu, \lambda) + |m_\varepsilon \circ \theta(\mu, \lambda)|^2) \mathcal{F}(f)(\mu, \lambda) \overline{\mathcal{F}(g)(\mu, \lambda)} d\gamma_{n+1}(\mu, \lambda), \quad (4.3)$$

and the norm $\|f\|_{\sigma,\rho} = \sqrt{\langle f, f \rangle_{\sigma,\rho}}$.

Lemma 4.4. Let $(s, y) \in [0, +\infty[\times \mathbb{R}^n$. Then

(i) The function

$$\Lambda_{(s,y)} : (\mu, \lambda) \mapsto \frac{\Phi_{(\mu,\lambda)}(s, y)}{\rho \sigma(\mu, \lambda) + |m_\varepsilon \circ \theta(\mu, \lambda)|^2},$$

belongs to $L^1(d\gamma_{n+1}) \cap L^2(d\gamma_{n+1})$.

(ii) The function

$$\Phi_{(s,y)} : (\mu, \lambda) \mapsto \frac{m_\varepsilon \circ \theta(\mu, \lambda) \Phi_{(\mu,\lambda)}(s, y)}{\rho \sigma(\mu, \lambda) + |m_\varepsilon \circ \theta(\mu, \lambda)|^2},$$

belongs to $L^1(d\gamma_{n+1}) \cap L^2(d\gamma_{n+1})$.

Where $\varphi_{(\mu,\lambda)}$ is the function given by relation (2.2).

Proof. The proof of the Lemma follows from relations (2.4), (4.1) and (4.2). \square

Proposition 4.5. Let $m \in L^\infty(d\nu_{n+1})$. Then the Hilbert space

$(\Omega_\sigma([0, +\infty[\times \mathbb{R}^n), \langle \cdot, \cdot \rangle_{\sigma,\rho})$ has the following reproducing Kernel

$$K_{\sigma,\rho}((r, x), (s, y)) = \int \int_{\Upsilon_+} \frac{\overline{\Phi_{(\mu,\lambda)}(r, x)} \Phi_{(\mu,\lambda)}(s, y)}{\rho \sigma(\mu, \lambda) + |m_\varepsilon \circ \theta(\mu, \lambda)|^2} d\gamma_{n+1}(\mu, \lambda), \quad (4.4)$$

that is

(i) For every $(s, y) \in [0, +\infty[\times \mathbb{R}^n$, the function $(r, x) \mapsto K_{\sigma,\rho}((r, x), (s, y))$ belongs to $\Omega_\sigma([0, +\infty[\times \mathbb{R}^n)$.

(ii) For every $f \in \Omega_\sigma([0, +\infty[\times \mathbb{R}^n)$, and $(s, y) \in [0, +\infty[\times \mathbb{R}^n$, we have the reproducing property,

$$\langle f, K_{\sigma,\rho}((\cdot, \cdot), (s, y)) \rangle_{\sigma,\rho} = f(s, y).$$

Proof. From Lemma 4.4 (i), the function $K_{\sigma,\rho}$ is well defined and by Theorem 2.3, we have

$$K_{\sigma,\rho}((r, x), (s, y)) = \mathcal{F}^{-1}(\Lambda_{(s,y)})(r, x), \quad (r, x) \in [0, +\infty[\times \mathbb{R}^n.$$

By Theorem 2.4, it follows that the function $K_{\sigma,\rho}((\cdot, \cdot), (s, y))$, belongs to $L^2(d\nu_{n+1})$, and we have

$$\mathcal{F}(K_{\sigma,\rho}((\cdot, \cdot), (s, y)))(\mu, \lambda) = \Lambda_{(s,y)}(\mu, \lambda), \quad (\mu, \lambda) \in \Upsilon. \quad (4.5)$$

Then by relations (2.4), (4.2) and (4.5), we obtain

$$\|K_{\sigma,\rho}((\cdot, \cdot), (s, y))\|_\sigma^2 \leq \frac{1}{\rho^2} \left\| \frac{1}{\sigma} \right\|_{1, \gamma_{n+1}}.$$

This proves that for every $(s, y) \in [0, +\infty[\times \mathbb{R}^n$, the function $K_{\sigma, \rho}((\cdot, \cdot), (s, y))$ belongs to $\Omega_{\sigma}([0, +\infty[\times \mathbb{R}^n)$.

(ii) From (4.3) and (4.5), we obtain

$$\langle f, K_{\sigma, \rho}((\cdot, \cdot), (s, y)) \rangle_{\sigma, \rho} = \iint_{\Upsilon_+} \mathcal{F}(f)(\mu, \lambda) \overline{\varphi_{(\mu, \lambda)}(s, y)} d\gamma_{n+1}(\mu, \lambda).$$

On the other hand, from relation (4.2) the function $\frac{1}{\sqrt{\sigma}}$ belongs to $L^2(d\gamma_{n+1})$, hence for every $f \in \Omega_{\sigma}([0, +\infty[\times \mathbb{R}^n)$, the function $\mathcal{F}(f)$ belongs to $L^1(d\gamma_{n+1})$. From this result and Theorem 2.3, we obtain

$$\langle f, K_{\sigma, \rho}((\cdot, \cdot), (s, y)) \rangle_{\sigma, \rho} = f(s, y).$$

This completes the proof of the Proposition. \square

Theorem 4.6. *Let $m \in L^{\infty}(d\nu_{n+1})$ and $\varepsilon > 0$, for every $h \in L^2(d\nu_{n+1})$ and $\rho > 0$, there exists a unique function $f_{\rho, h, \varepsilon}^*$ where the infimum*

$$\inf_{f \in \Omega_{\sigma}} \{ \rho \|f\|_{\sigma}^2 + \|h - T_{m, \varepsilon} f\|_{2, \nu_{n+1}}^2 \}, \quad (4.6)$$

is attained. Moreover the extremal function $f_{\rho, h, \varepsilon}^$ is given by*

$$f_{\rho, h, \varepsilon}^*(s, y) = \int_0^{+\infty} \int_{\mathbb{R}^n} h(r, x) \overline{V_{\sigma, \rho}((r, x), (s, y))} d\nu_{n+1}(r, x), \quad (4.7)$$

where $V_{\sigma, \rho}((r, x), (s, y)) = \iint_{\Upsilon_+} \frac{m_{\varepsilon} \circ \theta(\mu, \lambda) \overline{\varphi_{(\mu, \lambda)}(r, x)} \varphi_{(\mu, \lambda)}(s, y)}{\rho \sigma(\mu, \lambda) + |m_{\varepsilon} \circ \theta(\mu, \lambda)|^2} d\gamma_{n+1}(\mu, \lambda)$.

Proof. The existence and unicity of the extremal function $f_{\rho, h, \varepsilon}^*$ satisfying relation (4.6) is given by [10, 12, 21]. On the other hand From Proposition 4.2 and 4.5, we have

$$f_{\rho, h, \varepsilon}^*(s, y) = \langle h, T_{m, \varepsilon}(K_{\sigma, \rho})((\cdot, \cdot)(s, y)) \rangle_{\nu_{n+1}}, \quad (4.8)$$

where $\langle \cdot, \cdot \rangle_{\nu_{n+1}}$ denoted the inner product of $L^2(d\nu_{n+1})$, and $K_{\sigma, \rho}$ is the Kernel given by relation (4.4). According to Proposition 3.2 (ii), (4.5) and (4.8), we obtain

$$V_{\sigma, \rho}((r, x), (s, y)) = \iint_{\Upsilon_+} \frac{m_{\varepsilon} \circ \theta(\mu, \lambda) \overline{\varphi_{(\mu, \lambda)}(r, x)} \varphi_{(\mu, \lambda)}(s, y)}{\rho \sigma(\mu, \lambda) + |m_{\varepsilon} \circ \theta(\mu, \lambda)|^2} d\gamma_{n+1}(\mu, \lambda). \quad \square$$

Theorem 4.7. *Let $m \in L^{\infty}(d\nu_{n+1})$ and $h \in L^2(d\nu_{n+1})$. The extremal function $f_{\rho, h, \varepsilon}^*$ belongs to $\Omega_{\sigma}([0, +\infty[\times \mathbb{R}^n)$, and we have*

$$\|f_{\rho, h, \varepsilon}^*\|_{\sigma}^2 \leq \frac{1}{4\rho} \|h\|_{2, \nu_{n+1}}^2.$$

Proof. Let $(s, y) \in [0, +\infty[\times \mathbb{R}^n$. From Lemma 4.4 (ii) and Theorem 2.3, we have

$$V_{\sigma, \rho}((r, x), (s, y)) = \mathcal{F}^{-1}(\Phi_{(s, y)})(r, x).$$

By Theorem 2.4, it follows that the function $V_{\sigma, \rho}((\cdot, \cdot), (s, y))$ belongs to $L^2(d\nu_{n+1})$ and using Corollary 2.5, we get

$$\begin{aligned} f_{\rho, h, \varepsilon}^*(s, y) &= \int \int_{\Upsilon_+} \mathcal{F}(h)(\mu, \lambda) \overline{\Phi_{(s, y)}(\mu, \lambda)} d\gamma_{n+1}(\mu, \lambda) \\ &= \int \int_{\Upsilon_+} \mathcal{F}(h)(\mu, \lambda) \frac{\overline{m_\varepsilon \circ \theta(\mu, \lambda) \varphi_{(\mu, \lambda)}(s, y)}}{\rho \sigma(\mu, \lambda) + |m_\varepsilon \circ \theta(\mu, \lambda)|^2} d\gamma_{n+1}(\mu, \lambda). \end{aligned}$$

On the other hand, the function $(\mu, \lambda) \mapsto \mathcal{F}(h)(\mu, \lambda) \frac{\overline{m_\varepsilon \circ \theta(\mu, \lambda)}}{\rho \sigma(\mu, \lambda) + |m_\varepsilon \circ \theta(\mu, \lambda)|^2}$ belongs to $L^1(d\gamma_{n+1}) \cap L^2(d\gamma_{n+1})$, then by Theorem 2.3, we have

$$f_{\rho, h, \varepsilon}^*(s, y) = \mathcal{F}^{-1} \left(\mathcal{F}(h) \frac{\overline{m_\varepsilon \circ \theta(\cdot, \cdot)}}{\rho \sigma(\cdot, \cdot) + |m_\varepsilon \circ \theta(\cdot, \cdot)|^2} \right) (s, y).$$

From Theorem 2.4, it follows that, the function $f_{\rho, h, \varepsilon}^*$ belongs to $L^2(d\nu_{n+1})$, and we have for every $(\mu, \lambda) \in \Upsilon$,

$$\begin{aligned} \left| \mathcal{F}(f_{\rho, h, \varepsilon}^*)(\mu, \lambda) \right|^2 &= \left| \mathcal{F}(h)(\mu, \lambda) \frac{\overline{m_\varepsilon \circ \theta(\mu, \lambda)}}{\rho \sigma(\mu, \lambda) + |m_\varepsilon \circ \theta(\mu, \lambda)|^2} \right|^2, \quad (4.9) \\ &\leq \frac{1}{4\rho \sigma(\mu, \lambda)} |\mathcal{F}(h)(\mu, \lambda)|^2, \end{aligned}$$

thus, from Theorem 2.4 and Definition 4.1, we obtain

$$\|f_{\rho, h, \varepsilon}^*\|_\sigma^2 \leq \frac{1}{4\rho} \|h\|_{2, \nu_{n+1}}^2.$$

□

Theorem 4.8. Third Calderón's formula *Let $m \in L^\infty(d\nu_{n+1})$, and $f \in \Omega_\sigma([0, +\infty[\times \mathbb{R}^n)$. The extremal function $f_{\rho, \varepsilon}^*$ given by*

$$f_{\rho, \varepsilon}^*(s, y) = \int_0^{+\infty} \int_{\mathbb{R}^n} T_{m, \varepsilon} f(r, x) \overline{V_{\sigma, \rho}((r, x), (s, y))} d\nu_{n+1}(r, x),$$

satisfies

(i)

$$\lim_{\rho \rightarrow 0^+} \|f_{\rho, \varepsilon}^* - f\|_\sigma = 0. \quad (4.10)$$

(ii)

$$\lim_{\rho \rightarrow 0^+} f_{\rho, \varepsilon}^* = f, \quad \text{uniformly.}$$

Proof. (i) Let $f \in \Omega_\sigma([0, +\infty[\times \mathbb{R}^n)$, $h = T_{m,\varepsilon}f$, and $f_{\rho,\varepsilon}^* = f_{\rho,h,\varepsilon}^*$. From Proposition 4.2, the function h belongs to $L^2(d\nu_{n+1})$. Applying Definition 3.1, and relation (4.9), we obtain

$$\mathcal{F}(f_{\rho,\varepsilon}^*)(\mu, \lambda) = \mathcal{F}(f)(\mu, \lambda) \frac{|m_\varepsilon \circ \theta(\mu, \lambda)|^2}{\rho\sigma(\mu, \lambda) + |m_\varepsilon \circ \theta(\mu, \lambda)|^2}$$

Thus, it follows that for every $(\mu, \lambda) \in \Upsilon$

$$\mathcal{F}(f_{\rho,\varepsilon}^* - f)(\mu, \lambda) = \frac{-\rho\sigma(\mu, \lambda)\mathcal{F}(f)(\mu, \lambda)}{\rho\sigma(\mu, \lambda) + |m_\varepsilon \circ \theta(\mu, \lambda)|^2}. \quad (4.11)$$

Consequently,

$$\|f_{\rho,\varepsilon}^* - f\|_\sigma^2 = \iint_{\Upsilon_+} \frac{\rho^2\sigma^3(\mu, \lambda)|\mathcal{F}(f)(\mu, \lambda)|^2}{(\rho\sigma(\mu, \lambda) + |m_\varepsilon \circ \theta(\mu, \lambda)|^2)^2} d\gamma_{n+1}(\mu, \lambda).$$

Then, the result follows from the fact

$$\frac{\rho^2\sigma^3(\mu, \lambda)|\mathcal{F}(f)(\mu, \lambda)|^2}{(\rho\sigma(\mu, \lambda) + |m_\varepsilon \circ \theta(\mu, \lambda)|^2)^2} \leq \sigma(\mu, \lambda)|\mathcal{F}(f)(\mu, \lambda)|^2,$$

and the dominated convergence theorem.

(ii) By relation (4.2), the function $\frac{1}{\sqrt{\sigma}}$ belongs to $L^2(d\gamma_{n+1})$, hence for $f \in \Omega_\sigma([0, +\infty[\times \mathbb{R}^n)$, the function $\mathcal{F}(f)$ belongs to $L^1(d\gamma_{n+1})$. Then, from (4.11) and Theorem 2.3, we get

$$f_{\rho,\varepsilon}^*(s, y) - f(s, y) = \iint_{\Upsilon_+} \frac{-\rho\sigma(\mu, \lambda)\mathcal{F}(f)(\mu, \lambda)}{\rho\sigma(\mu, \lambda) + |m_\varepsilon \circ \theta(\mu, \lambda)|^2} \overline{\varphi_{(\mu, \lambda)}(s, y)} d\gamma_{n+1}(\mu, \lambda).$$

By using the dominated convergence theorem and the fact

$$\left| \frac{-\rho\sigma(\mu, \lambda)\mathcal{F}(f)(\mu, \lambda)\overline{\varphi_{(\mu, \lambda)}(s, y)}}{\rho\sigma(\mu, \lambda) + |m_\varepsilon \circ \theta(\mu, \lambda)|^2} \right| \leq |\mathcal{F}(f)(\mu, \lambda)|,$$

we deduce that

$$\lim_{\rho \rightarrow 0^+} \sup_{(s, y) \in [0, +\infty[\times \mathbb{R}^n} |f_{\rho,\varepsilon}^*(s, y) - f(s, y)| = 0.$$

Which completes the proof of the Theorem. \square

REFERENCES

- [1] G. Andrews, R. Askey and R. Roy, *Special functions*, Cambridge Univ. Press., New York 1999.
- [2] C. Baccar, S. Omri and L. T. Rachdi, *Fock spaces connected with spherical mean operator and associated operators*, *Mediterr. J. Math.*, **6** (2009), no. 1, 1–25.
- [3] M. Dziri, M. Jelassi and L. T. Rachdi, *Spaces of DL_p type and a convolution product associated with the spherical mean operator*, *Int. J. Math. Math. Sci.*, **2005** (2005), no. 3, 357–381.
- [4] J. A. Fawcett, *Inversion of N -dimensional spherical means*, *SIAM. J. Appl. Math.*, **45** (1985), 336–341.
- [5] J. Gosselin and K. Stempak, *A weak-type estimate for Fourier-Bessel multipliers*, *Proc. of the Amer. Math. Soc.*, vol**106**,no. 3 (1989), 655–662.
- [6] H. Helesten and L. E. Anderson, *An inverse method for the processing of synthetic aperture radar data*, *Inverse Problems.*, **4** (1987), 111–124.
- [7] M. Herberthson, *A numerical implementation of an inverse formula for CARABAS raw Data.*, Internal Report D 30430-3.2, National Defense Research Institute, FOA, Box 1165; S-581 11, Linköping, Sweden, 1986.
- [8] M. Jelassi and L. T. Rachdi, *On the range of the Fourier transform associated with the spherical mean operator*, *Fract. Calc. Appl. Anal.*, **7** (2004), no. 4, 379–402.
- [9] Kapelko, R.: *A multiplier theorem for the Hankel transform*. *Rev. Mat. Complut.* **11**, 281288 (1998).
- [10] G.S. Kimeldorf and G.Wahba, *Some results on Tchebycheffian spline functions*, *J. Math. Anal. Appl.* **33** (1971), 82–95.
- [11] N. N. Lebedev, *Special functions and their applications*, Dover publications, New York 1972.
- [12] T. Matsuura, S. Saitoh and D.D. Trong, *inversion formulas in heat conduction multidimensional spaces*, *J. Inverse III-posed Problems* **13** (2005), 479–493.
- [13] N. Msehli and L.T. Rachdi, *Beurling-Hörmander uncertainty principle for the spherical mean operator*, *J. Ineq. Pure and Appl. Math.*, **10** (2009), Iss. 2, Art. 38.
- [14] N. Msehli and L.T. Rachdi, *Heisenberg-Pauli-Weyl uncertainty principle for the spherical mean operator*, *Mediterr. J. Math.*, **7** (2010), no. 2, 169–194.
- [15] M. M. Nessibi, L. T. Rachdi, and K. Trimèche, *Ranges and inversion formulas for spherical mean operator and its dual*, *J. Math. Anal. Appl.*, **196** (1995), no. 3, 861–884.
- [16] S. Omri, *Uncertainty principle in terms of entropy for the spherical mean operator*, *J. Math. Ineq.*, **5** (2011), Issue 4, 473–490.
- [17] L. Peng and J. Zhao, *Wavelet and Weyl transforms associated with the spherical mean operator*, *Integr. equ. oper. theory.*, **50** (2004), 279-290.
- [18] L. T. Rachdi, and K. Trimèche, *Weyl transforms associated with the Spherical Mean Operator*, *Anal. Appl.*, **1** (2003), N2, 141–164.
- [19] S. Saitoh, *Hilbert spaces induced by Hilbert space valued functions*, *Proc. Amer.*

- Math. Soc. **89** (1983), 74-78.
- [20] S. Saitoh, *The Weierstrass transform and an isometry in the heat equation*, Appl. Anal. **16** (1983) 1-6.
- [21] S. Saitoh, *Approximate real inversion formulas of the Gaussian convolution*, Appl. Anal. **83**, (2004), 727-733.
- [22] S. Saitoh, *Best approximation, Tikhonov regularization and reproducing kernels*, Kodai Math. J. **28** (2005) 359-367.
- [23] T. Schuster, *The method of approximate inverse: theory and applications*, Lecture Notes in Math, Vol. **1906**, 2007.
- [24] F. Soltani, *L^p - Fourier multipliers for the Dunkl operator on the real line*, J. Functional Anal. **209** (2004), 16-35.
- [25] F. Soltani, "Multiplier operators and extremal functions related to the dual Dunkel-Sonine operator", Acta Mathematica Scientia. Series B: English Edition, Vol. **33**, no. 2, (2013), 430-442.
- [26] K. Trimèche, *Transformation intégrale de Weyl et théorème de Paley-Wiener associés à un opérateur différentiel singulier sur $(0, \infty)$* , J. Math. Pures et Appl. **60** (1981), 51-98.
- [27] K. Trimèche, *Inversion of the Lions translation operator using generalized wavelets*, Appl. Comput. Harmonic Anal. **4** (1997), 97-112.
- [28] L. V. Wang, *Photoacoustic imaging and spectroscopy*, Optical science and engineering, Vol. **144**, 2009.
- [29] G. N. Watson, *A treatise on the theory of Bessel functions*, 2nd ed., Cambridge Univ. Press., London/New York, (1966).
- [30] J. Zhao, *Localization operators associated with the spherical mean operator*, Anal. Theory Appl., **21** (2005), no. 4, 317-325.

khaled.hleili@gmail.com

Department of Mathematics, Faculty of Science, Northern Borders University,
Arar, Saudi Arabia

Preparatory Institute for Engineering Studies of Kairouan

Department of Mathematics

Avenue Assad Ibn Fourat Kairouan 3100

KHALED HLEILI

Department of Mathematics, Faculty of Science, Northern Borders University,
Arar, Saudi Arabia

Preparatory Institute for Engineering Studies of Kairouan

Department of Mathematics

Avenue Assad Ibn Fourat Kairouan 3100

e-mail: khaled.hleili@gmail.com